

Acute leukemias

**Draft guide to the completion of the EBMT data collection
form: Acute_Leukemias_v2.0**

May 2024

EBMT Registry

EBMT Clinical Research & Registry Department



**Co-funded by
the European Union**

Co-funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or European Health and Digital Executive Agency (HADEA). Neither the European Union nor the granting authority can be held responsible for them.

Table of Contents

Introduction	5
Acute leukaemias	5
Disease	5
Date of diagnosis	5
Acute leukaemias main classification	5
Haematological values	6
Peripheral blood	6
White Blood cell count (10 ⁹ cells/L)	6
% of blasts	6
Bone marrow	6
% of blasts	6
Acute myeloid leukaemias (AML)	6
Disease	6
Classification	6
AML with myelodysplasia related changes?	6
Was there a previous diagnosis of MDS, MPN or MDS/MPN?	7
Therapy-related myeloid neoplasia (old "secondary acute leukaemia")	7
Chromosome analysis	7
Chromosome analysis done at diagnosis	7
Date of chromosome analysis	8
What were the results?	8
Number of abnormalities	8
Complex karyotype	8
Monosomal karyotype	8
Multiple trisomies	9
Chromosome analysis details	9
Transcribe the complete karyotype	10
Molecular marker analysis	10
Molecular marker analysis at diagnosis	10
Date of molecular markers analysis	11
Molecular marker analysis details	11
Next generation sequencing (NGS) performed at diagnosis	13
Disease	13
Other AML classification	13
FAB classification of AML	13
Involvement at time of diagnosis	14
Medullary involvement	14
Extramedullary involvement	14
Organs involved at time of diagnosis	14

Precursor lymphoid neoplasms (previously ALL)	14
Disease	14
Classification	14
Secondary origin: is this PLN related to prior exposure to therapeutic drugs or radiation?	15
Due to exposure to	15
Chromosome analysis	15
Chromosome analysis done at diagnosis	15
Date of chromosome analysis	16
What were the results?	16
Number of abnormalities	16
Complex karyotype	16
Chromosome analysis details	16
Transcribe the complete karyotype	17
Molecular marker analysis	17
Molecular marker analysis at diagnosis	17
Date of molecular markers analysis	17
Molecular marker analysis details	18
Ph-like ALL?	19
Next generation sequencing (NGS) performed at diagnosis	20
Disease	20
Involvement at time of diagnosis	20
Medullary involvement	20
Extramedullary involvement	20
Organs involved at time of diagnosis	20
Other acute leukaemias	21
Disease	21
Classification	21
Secondary origin: is this other acute leukaemia related to prior exposure to therapeutic drugs or radiation?	21
Due to exposure to	21
Chromosome analysis	22
Chromosome analysis at diagnosis	22
Date of chromosome analysis	22
What were the results?	22
Number of abnormalities	22
Complex karyotype	23
Chromosomal abnormalities	23
Transcribe the complete karyotype	23
Involvement at time of diagnosis	23
Medullary involvement	23
Extramedullary involvement	23

Disease	23
Organs involved at time of diagnosis	23
Bibliography	25

Introduction

Please make sure you have already checked the **Introduction to the EBMT Registry Completion Guidelines** document latest version available under *Manuals and Reference Documents* section on [EBMT website](#).

Acute leukaemias

Acute leukaemia is a malignant disease that originates either in a lymphopoietic stem cell (Precursor lymphoid neoplasms (PLN), previously ALL) or in a hematopoietic stem cell or progenitor cell (acute myeloid leukaemia, AML). Other Acute Leukaemias with ambiguous lineage can be reported with this form.

Complete this form only if this diagnosis was the indication for a HCT/CT/GT or if it was specifically requested.

Disease

Date of diagnosis

Report the date of the first pathological diagnosis of the disease. Add the date when the sample was collected for examination or (in its absence) the date indicated by a physician within the patient's medical record.

Acute leukaemias main classification

Select the main class that is appropriate for acute leukaemia and check the box next to it.

There are distinguished 3 main classes of acute leukaemia, which are:

- **Acute myeloid leukaemia (AML):** characterised by disordered differentiation and proliferation of hematopoietic stem cells or progenitor cells into myeloblasts in acute myeloid leukaemia.
- **Precursor lymphoid neoplasms (PLN, previously ALL)** characterised by disordered differentiation and proliferation of lymphopoietic cells into lymphoblasts in precursor lymphoid neoplasms.
- **Other acute leukaemia** of ambiguous lineage: Acute undifferentiated leukaemia, Mixed phenotype, Natural killer (NK) - cell lymphoblastic leukaemia/lymphoma or Other leukaemia.

Haematological values

Peripheral blood

Report the White Blood cells (WBC) count at diagnosis in peripheral blood. The white blood cell count is expressed in two ways: as an absolute number and as a percentage.

White Blood cell count (10^9 cells/L)

Indicate the absolute number of white blood cells at diagnosis or mark it as **Not evaluated** or **Unknown**.

Please, pay attention to the units for the number of WBC reported at diagnosis: 10^9 cells/L is the same as 10^6 cells/ml or 10^3 cells/ μ l. If the WBC count is given in mm^3 then it should be transformed and reported in 10^9 cells/L: 300.000mm^3 is equal to 300×10^9 cells/L.

% of blasts

Indicate the percentage of blasts in peripheral blood at diagnosis. The percentage of blasts can be found in the white blood cell count.

Bone marrow

% of blasts

Indicate the percentage of blasts in bone marrow at diagnosis. It can be found in a bone marrow aspirate laboratory result (not in the bone marrow “histology / biopsy” form). If this data is not available, mark it as **Not evaluated** or **Unknown**, whatever is more appropriate.

Acute myeloid leukaemias (AML)

Disease

Classification

AML with myelodysplasia related changes?

In most cases, this classification applies to AML where an MDS, MPN or an MDS/MPN has been diagnosed beforehand. In a few cases, it applies to what looks as a de novo AML at the time of diagnosis, but which after further analysis of the bone marrow or after treatment for the AML, there is a suggestion that there could have been an undetected history of myelodysplastic syndrome (MDS).

Indicate whether AML with myelodysplasia related changes was diagnosed or not, tick the box **Unknown** if it is not known. If the answer is **Yes**, specify also if the patient was previously diagnosed with MDS or MDS/MPN in the subquestion.

Was there a previous diagnosis of MDS, MPN or MDS/MPN?

If the patient had a previous diagnosis of MDS, MPN or MDS/MPN, answer **Yes** and fill-in and submit respective indication diagnosis form in addition to the current form; otherwise, answer **No**. If it is unknown whether or not the patient had a previous diagnosis of MDS, MPN or MDS/MPN, check **Unknown**.

Note: For AML with myelodysplasia related changes, if MDS, MPN or MDS/MPN was previously diagnosed, besides the Acute leukaemia form the corresponding indication diagnosis form for MDS, MPN or MDS/MPN must be filled in as well.

Therapy-related myeloid neoplasia (old "secondary acute leukaemia")

Indicate whether therapy-related myeloid neoplasia (therapy-related myeloid neoplasms or t-MN) was diagnosed or not, tick the box **Unknown** if it is not known.

Therapy-related myeloid neoplasia arises as a late-effect of cytotoxic and/or radiation therapy for a primary diagnosis. Answer **Yes** if it is related to prior treatment, but not after a previous diagnosis of MDS, MPN or MDS/MPN.

- If the patient has therapy-related AML and did not undergo HCT/CT for the previous diagnosis, please complete the *non-indication diagnosis* form to record the previous diagnosis.
- If the patient has AML related to HCT/CT therapy treatment for the previous diagnosis, please complete the appropriate *indication diagnosis* form for this previous diagnosis.

Chromosome analysis

In this section describe the results of the most complete chromosome analysis performed around the time of diagnosis.

Chromosome analysis done at diagnosis

Report if the chromosome analysis has been done around the time of diagnosis or not. Choose the answer based on the following:

No - the chromosome analysis has not been done or failed;

Yes - the chromosome analysis has been performed for diagnosis, specify also the **Output of analysis**: if it is **Separate abnormalities** or **Full karyotype**;

Unknown - it is unknown whether the chromosome analysis has been done or not.

Date of chromosome analysis

If chromosome analyses was done around the time of diagnosis, please specify the date of the chromosome analyses or mark it as **Unknown**.

What were the results?

Indicate the results of the chromosome analysis you are reporting and choose the answer based on the following:

Normal results - the chromosome analysis has been performed and all the results have been found normal (46,XY or 46,XX).

Abnormal results - the chromosome analysis has been performed and at least one of the results has been found to be abnormal. Provide further details on abnormal results in the sub questions below.

Failed - the chromosome analysis has failed.

Number of abnormalities

If the results were **abnormal**, indicate the number of abnormalities present in the most complete analysis with abnormal results.

Complex karyotype

A complex karyotype is defined by ≥ 3 unrelated chromosome abnormalities in the absence of other class-defining recurrent genetic abnormalities; excludes hyperdiploid karyotypes with three or more trisomies (or polysomies) without structural abnormalities.

Indicate whether the karyotype is complex or not, check the corresponding **Unknown** box if it is not known.

Monosomal karyotype

A monosomal karyotype is defined by the presence of two or more distinct monosomies (excluding loss of X or Y), or one single autosomal monosomy in combination with at least one structural chromosome abnormality (excluding core-binding factor AML).

Indicate whether it is monosomal karyotype or not, check the corresponding **Unknown** box if it is not known.

Multiple trisomies

If there is more than one trisomy please answer **Yes**, otherwise answer **No**. Check the corresponding **Unknown** box if it is not known.

Chromosome analysis details

See the cytogenetics form or ask the cytogenetics team and consult your physician.

If chromosome analysis was performed and had abnormal results, indicate for each abnormality listed in the table whether it was **Absent** or **Present**. If a chromosome abnormality was not evaluated, report **Not evaluated**.

If a chromosome abnormality was checked, but not listed as an option in the table, select **Other** and specify the abnormality, marking whether it was **Absent** or **Present**.

If **11q23**, **3q26 (EVI1)**, **abn 5 type** or **abn 7 type** abnormality was present, mark for each subtype abnormality if it was **Absent**, **Present** or **Not evaluated**. If the subtype is not listed, please use corresponding Other option and specify it in the textfield and report if it was **Absent** or **Present**.

Please find below the minimum list of requested cytogenetics abnormalities according to the WHO classification 2022:

- t(15;17)
- t(8;21)
- inv(16)/ t(16;16)
- 11q23 abnormality type, if a 11q23 abnormality is present:
 - t(9;11)
 - t(11;19)
 - t(10;11)
 - t(6;11)
 - Other abn(11q23); specify.
- 3q26 (EVI1) abnormality type, if a 3q26 abnormality is present:
 - inv(3) / t(3;3)
 - t(2;3)(p21;q26)
 - Other (3q26)/EVI1 rearrangement; specify.
- t(6;9)
- abn 5 type, if an abn 5 is present:
 - del (5q)

- monosomy 5
 - add(5q)
 - Other abn(5q); specify.
- abn 7 type, if an abn 7 is present:
 - del(7q)
 - monosomy 7
 - add(7q)
 - Other abn(7q); specify.
- Monosomy 17
- abn(17p)
- t(1;22)
- Trisomy 8
- t(9;22)
- t(8;16)
- Other; specify.

Transcribe the complete karyotype

If it is not possible to report the chromosome analysis results as per abnormalities table above, please report the full complete karyotype.

Molecular marker analysis

Molecular marker analysis at diagnosis

Molecular markers are specific genetic sequences that are associated with the recipient's primary disease. These markers are determined by molecular biology laboratories that have expertise in this field. They are used for diagnosis and sometimes to follow up on the disease after treatment.

Indicate whether molecular biology studies have been done around the time of diagnosis to identify molecular markers. If they have been done, select **Yes**. If no molecular biology has been done, please check **No**. Select **Unknown** if it is unknown whether the analysis of the molecular markers has been done or not.

Date of molecular markers analysis

If molecular biology studies have been done around the time of diagnosis to identify molecular markers, please specify the date of the molecular markers analysis which results will be reported below. Select **Unknown** if the date is unavailable.

Molecular marker analysis details

If molecular marker analysis was performed, indicate for each marker in the table whether it was **Absent** or **Present**. If a molecular marker was not evaluated, report **Not evaluated**.

If a molecular marker was evaluated, but not listed as an option in the table, select **Other** and specify the marker, indicating whether it was **Absent** or **Present**.

See the list below:

- AML1-ETO (RUNX1/RUNX1T1) Molecular product of t(8;21)
- CBFβ-MYH11 Molecular product of inv(16)(p13.1;q22) or (16;16)(p13.1;q22)
- PML-RAR Molecular product of t(15;17)
- MLL (KMT2A)-rearrangement/mutation:
 - MLLT3(AF9)-MLL
 - Molecular product of t(9;11)(p22;q23)
 - MLL-PTD
 - (partial tandem duplication)
 - MLLT4(AF6)-MLL
 - Molecular product of t(6;11)(q27;q23)
 - ELL-MLL
 - Molecular product of t(11;19)(q23;p13.1)
 - MLLT1(ENL)-MLL
 - Molecular product of t(11;19)(q23;p13.3)
 - MLLT10(AF10)-MLL
 - Molecular product of t(10;11)(p12;q23)
 - Other MLL-rearrangement; specify:
- DEK-NUP214(CAN) Molecular product of translocation t(6;9)(p23;q34)
- RPN1-EVI1 Molecular product of inv(3)(q21q26.2) or t(3;3)(q21q26.2)
- RBM15-MKL1 Molecular product of translocation t(1;22)(p13;q13)
- NPM1 mutation
- c-KIT

- DNMT3A
- ASXL1
- TP53
- RUNX1
- IDH2
- IDH1
- BRAT
- SRSF2
- SF3B1
- CEBPA: if CEBPA present, specify also whether or not there was bZIP mutation (answer yes, no, unknown) and if the mutation is biallelic (answer yes, no, unknown).
- FLT3-ITD (internal tandem duplication)
- FLT3-TKD
- BCR-ABL
- GATA2
- MECOM(EVI1)
- KAT6A-CREBBP
- BCOR
- EZH2
- STAG2
- U2AF1
- ZRSR2
- Other; specify.

If **MLL** rearrangement/mutation was present, specify for each of its subtypes in the table if it was **Absent**, **Present** or **Not evaluated**; if there were checked other subtypes that are not mentioned in the table, specify the rearrangement/mutation in the **Other MLL-rearrangement** text field and if it was **Absent** or **Present**.

For **CEBPA mutation**, if present, please specify if

- there is a mutation in the CEBPA bZIP?
- the mutation is biallelic (either in bZIP or other mutation in the CEBPA gene) .

Next generation sequencing (NGS) performed at diagnosis

If NGS was done at diagnosis, select **Yes**. Otherwise, please check **No**. Select **Unknown** if it is unknown whether the NGS has been done or not.

Disease

Other AML classification

If applicable, select the corresponding WHO classification of other AML diagnosis (1):

- Acute panmyelosis with myelofibrosis
- Myeloid sarcoma (granulocytic sarcoma)
- Myeloid proliferations related to Down syndrome
- Blastic plasmacytoid dendritic cell neoplasm (BPDCN)

FAB classification of AML

If available, indicate the appropriate FAB class.

The French-American-British (FAB) classification for acute leukaemias is a classification system that relies on how the cells appear under a microscope. The following AML classes are distinguished (2):

FAB subtype	Name
M0	AML with minimal differentiation
M1	AML without maturation
M2	AML with maturation
M3	Acute promyelocytic leukaemia
M4	Acute myelomonocytic leukaemia
M5	Acute monoblastic and monocytic leukaemia
M6	Acute erythroid leukaemia
M7	Acute megakaryoblastic leukaemia

Table 1, FAB classification of AML (2)

Involvement at time of diagnosis

Medullary involvement

If the AML, indicate whether the Marrow / Blood was involved answering **yes**, **no** or **unknown**. Please be aware that although the vast majority of acute leukaemias involve the invasion of the bone marrow by blasts, there are cases where blast invasion is only found in organs other than the bone marrow (e.g. choromas).

Extramedullary involvement

Extramedullary involvement (EMI) refers to leukemic cells found in organs or tissue outside the blood or bone marrow. The most common sites of extramedullary disease are the central nervous system (CNS), skin and ovaries/testes.

Please indicate if any other organ(s) than the Marrow or Blood were involved at diagnosis, by reporting extramedullary involvement and answering **yes**, **no** or **unknown**.

Organs involved at time of diagnosis

Indicate per each organ in the list if leukaemic cells were found there (answer **Yes**) or not (answer **No**), or if it was **Not evaluated** at time of diagnosis. If there were checked organs other than from the list, check the **Other** box and specify the organ, indicating if it is involved (select **Yes**) or not (select **No**).

See the list below:

- Skin:
- CNS:
- Testes/Ovaries:
- Other; specify.

Precursor lymphoid neoplasms (previously ALL)

Disease

Classification

Select the class that is appropriate for the precursor lymphoid neoplasm (PLN) and check the box next to it. If the class is not listed, check the box **Other precursor lymphoid neoplasm** and specify it in the textbox. Select from the following answer options:

- B lymphoblastic leukaemia/lymphoma
- T lymphoblastic leukaemia/lymphoma
- Other precursor lymphoid neoplasm; specify.

Secondary origin: is this PLN related to prior exposure to therapeutic drugs or radiation?

Indicate if this precursor lymphoid neoplasm is related to prior exposure to therapeutic drugs or radiation, i.e. has a **secondary origin** answering **yes**, **no** or **unknown**. If the answer is **Yes**, indicate also:

Due to exposure to

Select what the patient was exposed to, that caused PLN. Check the respective check box if it was:

- Chemotherapy / radiotherapy treated disease
- Immune suppression
- Other; specify _____
- Unknown: if PLN is considered to be of secondary origin but the exact reason cannot be identified.

If the PLN is related to prior exposure to therapeutic drugs or radiation, please make sure that the diagnosis of the prior treatment is reported in the registry:

- if no HCT/CT was done for this previous diagnosis, please use a **Non-Indication Diagnosis** form, or
- if the patient had HCT/CT for the previous diagnosis, use the appropriate indication **diagnosis** form to report it, unless already registered in EBMT Registry.

Chromosome analysis

Chromosome analysis done at diagnosis

In this section describe the results of the most complete chromosome analysis performed around the time of diagnosis. Choose the answer based on the following:

No - the chromosome analysis has not been done.

Yes - the chromosome analysis has been performed for diagnosis.

Unknown - it is unknown whether the chromosome analysis has been done or not.

Date of chromosome analysis

If chromosome analysis was performed, please specify the date of the chromosome analysis around the time of diagnosis, which results will be reported. Mark this field as **unknown** if the date is not available.

What were the results?

Indicate the results of the chromosome analysis you are reporting and choose the answer based on the following:

Normal results - the chromosome analysis has been performed and all the results have been found normal (46,XY or 46,XX).

Abnormal results - the chromosome analysis has been performed and at least one of the results has been found to be abnormal.

Failed - the chromosome analysis has failed.

Number of abnormalities

- If the results of the chromosome analysis were **abnormal**, indicate the number of abnormalities present in the most complete analysis with abnormal results

Complex karyotype

A complex karyotype is defined by ≥ 3 unrelated chromosome abnormalities in the absence of other class-defining recurrent genetic abnormalities; excludes hyperdiploid karyotypes with three or more trisomies (or polysomies) without structural abnormalities.

If the results of the chromosome analysis were **abnormal**, indicate whether the karyotype is complex or not, check the corresponding **Unknown** box if it is not known.

Chromosome analysis details

See the cytogenetics form or ask the cytogenetics team and consult your physician.

If chromosome analysis was performed and had abnormal results, indicate for each abnormality listed in the table whether it was **Absent** or **Present**. If a chromosome abnormality was not evaluated, report **Not evaluated**.

If a chromosome abnormality was checked, but not listed as an option in the table, select **Other** and specify the abnormality, marking whether it was **Absent** or **Present**.

If **11q23** abnormality was present, specify also if t(4;11) was **Absent, Present** or **Not evaluated**. If another 11q23 sub-type was checked, specify it in the **Other abn(11q23)** text field and mark if it was **Absent** or **Present**.

If **hyperdiploidy (>46 chromosomes)** is present, specify also the following:

- If 51-67 chromosomes were **Absent, Present** or **Not evaluated** in the chromosome analysis
- In case of trisomy; specify the extra chromosome name and if it was **Absent, Present** or **Not evaluated**.
- If **other hyperdiploid karyotype**; specify the number of chromosomes in the textbox and if they were **Absent** or **Present**.

For hypodiploidy (<46 chromosomes), if hypodiploidy is present, answer subquestions to specify the number of missing chromosomes by marking if low hypodiploid (32 - 39 chromosomes) and near haploid (24-31 chromosomes), monosomy (specify details of monosomy in the text field) were **Absent, Present** or **Not evaluated** in the chromosome analysis. In case of other chromosome abnormality, specify the number of chromosomes in **Other; number of chromosomes** field and mark if they were **Absent** or **Present**.

Transcribe the complete karyotype

If it is not possible to report the chromosome analysis results as per abnormalities table above, please report the full complete karyotype .

Molecular marker analysis

Molecular marker analysis at diagnosis

Indicate whether molecular biology studies have been done to identify molecular markers. If they have been done, select **Yes**. If no molecular biology has been done, please check **No**. Select **Unknown** if it is unknown whether the analysis of the molecular markers has been done or not.

Date of molecular markers analysis

If molecular biology has been done, report the date of the molecular marker analysis. Select **Unknown** if the date is unavailable.

Molecular marker analysis details

If molecular marker analysis was performed, indicate for each marker in the table whether it was **Absent** or **Present**. If a molecular marker was not evaluated, report **Not evaluated**.

If a molecular marker was evaluated, but not listed as an option in the table, select **Other** and specify the marker, indicating whether it was **Absent** or **Present**.

For **MLL (KMT2A)** rearrangement/mutation, if it was present, specify for each subtype in the table if it was **Absent**, **Present** or **Not evaluated**; if there were checked other subtypes that are not mentioned in the table, specify the rearrangement/mutation in the **Other MLL (KMT2A)-rearrangement** text field and if it was **Absent** or **Present**.

Please see the list of molecular markers requested for PLN:

- BCR-ABL Molecular product of t(9;22)(q34;q11.2)
- PML-RAR Molecular product of t(15;17)
- MLL (KMT2A)-rearrangement/mutation:
 - AFF1(AF4)-MLL M
 - Molecular product of t(4;11)(q21;q23)
 - MLLT1(ENL)-MLL
 - Molecular product of t(11;19)(q23;p13.3)
 - MLLT3(AF9)-MLL
 - Molecular product of t(9;11)(p22;q23)
 - Other MLL-rearrangement; specify.
- TEL(ETV6)-AML1(RUNX1) Molecular product of t(12;21)(p13;q22)
- IL3-IGH Molecular product of translocation t(5;14)(q31;q32)
- TCF3-PBX1 Molecular product of translocation (1;19)(q23;p13.3)
- IKZF1 (IKAROS) deletion or mutation
- NOTCH1 / FBWX7 mutation
- PAX5 mutation
- KRAS mutation
- NRAS mutation
- PTEN mutation
- FLT3 mutation
- PTPN11 mutation
- BCL/MYC-rearranged
- Other; specify.

Ph-like ALL?

Philadelphia (Ph)-like acute lymphoblastic leukemia (ALL) is a high-risk subtype of B-precursor ALL (B-ALL) characterised by a gene-expression profile similar to Ph⁺ ALL but lacking the specific *BCR-ABL1* fusion gene. They could be treated by TKIs.

These alterations can be subdivided into 5 distinct subgroups based on the type of cytokine receptor or kinase fusion present:

(1) rearrangements of *CRLF2*: ex: *P2RY8-CRLF2* rearrangements, *IGH@-CRLF2*.

(2) ABL-class gene rearrangements, such as *ABL1*, *ABL2*, *PDGFRB*, and *CSF1R*

(3) *JAK2* and *EPOR* rearrangements.

(4) sequence mutations or deletions activating JAK-STAT- or MAPK signalling pathways. For example: mutations of *IL7R*, *FLT3*, and *IL2RB*, mutations of *JAK1* and *JAK3*, mutation of *SH2B3*.

(5) other rare kinase alterations (ex: mutations of *NRAS*, *KRAS*, *PTPN11*, and *NF1*).

Indicate if Philadelphia chromosome (Ph)-like acute lymphoblastic leukaemia (ALL) was analysed or not by answering:

- **No** in case it was not checked (skip the table below),
- **Yes** to mark it was investigated (proceed to report its results in the table below), or
- **Unknown** if it is not known whether such investigation was done or not.

If the (Ph)-like acute lymphoblastic leukaemia (ALL) was analysed, complete the table with alterations and mark per each rearrangement (alteration) if it was **Absent**, **Present** or **Not evaluated**.

Please see the list of molecular marker of interest for Ph-like:

- CRFL2-P2RY8
 - Other CRFL2 rearrangement; specify.
- ABL1 rearrangement:
 - ABL1-ETV6
 - ABL1-NUP214
 - Other ABL1 rearrangement; specify.
- ABL2 rearrangement:

- ABL2-RCSD1
 - Other ABL2 rearrangement; specify.
- JAK2 rearrangement:
 - JAK2-PAX5
 - JAK2-BCR
 - Other JAK2 rearrangement; specify.
- EPOR rearrangement:
 - EPOR-IGH
 - Other EPOR rearrangement; specify.

Next generation sequencing (NGS) performed at diagnosis

If NGS was done, select **Yes**. Otherwise, please check **No**. Select **Unknown** if it is unknown whether the NGS has been done or not.

Disease

Involvement at time of diagnosis

Medullary involvement

Indicate whether the Marrow / Blood was involved answering **yes**, **no** or **unknown**. Please be aware that although the vast majority of acute leukaemias involve the invasion of the bone marrow by blasts, there are cases where blast invasion is only found in organs other than the bone marrow (e.g. choromas).

Extramedullary involvement

Extramedullary involvement (EMI) refers to leukemic cells found in organs or tissue outside the blood or bone marrow. The most common sites of extramedullary disease are the central nervous system (CNS), skin and ovaries/testes.

Please indicate if any other organ(s) than the Marrow or Blood were involved at diagnosis, by reporting extramedullary involvement and answering **yes**, **no** or **unknown**.

Organs involved at time of diagnosis

Indicate per each organ in the list if leukemic cells were found there (answer **Yes**) or not (answer **No**), or if it was **Not evaluated** at time of diagnosis. If there were checked organs other than from the list, check the **Other** box and specify the organ, indicating if it is involved (select **Yes**) or not (select **No**).

See the list below:

- Skin;
- CNS;
- Testes/Ovaries;
- Other; specify.

Other acute leukaemias

Disease

Classification

Acute leukaemias of ambiguous lineage may be divided into:

Acute undifferentiated leukaemia (AUL). AULs are leukaemias with very primitive phenotypes with little evidence of lineage commitment.

Mixed-phenotype acute leukaemias (MPALs). MPAL is a heterogeneous category that encompasses rare blastic hematopoietic cell neoplasms that express a mixture of myeloid and lymphoid (B- or T-lineage) antigens.

Natural killer (NK) - cell lymphoblastic leukaemia/lymphoma.

Indicate the class that is appropriate for acute leukaemia of ambiguous lineage. If the class is not listed, tick the box **Other** and specify it in the text field.

Secondary origin: is this other acute leukaemia related to prior exposure to therapeutic drugs or radiation?

Indicate if this acute leukaemia is related to prior exposure to therapeutic drugs or radiation, i.e. has a secondary origin. If the answer is **Yes**, indicate also:

Due to exposure to

What the patient was exposed to, that caused this acute leukaemia. Check the respective check box if it was chemotherapy/radiotherapy treated disease or immune suppression, otherwise check the **Other** box and specify. Select **Unknown** if this acute leukaemia is considered to be of secondary origin but the exact reason cannot be identified.

Note: If not reported yet, complete and submit the respective **non-indication diagnosis** form in addition to the current form to report the diagnosis in regard to which drugs/therapy that caused acute leukaemia were applied.

Chromosome analysis

Chromosome analysis at diagnosis

In this section describe the results of the most complete chromosome analysis performed around the time of diagnosis.

Choose the answer based on the following:

No - the chromosome analysis has not been done.

Yes - the chromosome analysis has been performed for diagnosis.

Unknown - it is unknown whether the chromosome analysis has been done or not.

Date of chromosome analysis

If chromosome analysis was performed around the time of diagnosis, please specify the date of the chromosome analysis.

What were the results?

Indicate the results of the chromosome analysis you are reporting and choose the answer based on the following:

Normal results - the chromosome analysis has been performed and all the results have been found normal (46,XY or 46,XX).

Abnormal results - the chromosome analysis has been performed and at least one of the results has been found to be abnormal.

Failed - the chromosome analysis has failed.

Number of abnormalities

If the results were **abnormal**, indicate the number of abnormalities present in the most complete analysis with abnormal results.

Complex karyotype

A complex karyotype is defined by ≥ 3 unrelated chromosome abnormalities in the absence of other class-defining recurrent genetic abnormalities; excludes hyperdiploid karyotypes with three or more trisomies (or polysomies) without structural abnormalities.

Indicate whether the karyotype is complex or not, check the corresponding **Unknown** box if it is not known.

Chromosomal abnormalities

If the results of the chromosome analysis were **abnormal**, specify the chromosome abnormalities in the text field and mark whether they were **Absent** or **Present**.

Transcribe the complete karyotype

If it is not possible to report the chromosome analysis results as per abnormalities table above, please report the full complete karyotype.

Involvement at time of diagnosis

Medullary involvement

Indicate whether the Marrow / Blood was involved answering **yes**, **no** or **unknown**. Please be aware that although the vast majority of acute leukaemias involve the invasion of the bone marrow by blasts, there are cases where blast invasion is only found in organs other than the bone marrow (e.g. choromas).

Extramedullary involvement

Extramedullary involvement (EMI) refers to leukemic cells found in organs or tissue outside the blood or bone marrow. The most common sites of extramedullary disease are the central nervous system (CNS), skin and ovaries/testes.

Please indicate if any other organ(s) than the Marrow or Blood were involved at diagnosis, by reporting extramedullary involvement and answering **yes**, **no** or **unknown**.

Disease

Organs involved at time of diagnosis

Indicate per each organ in the list if leukemic cells were found there (answer **Yes**) or not (answer **No**), or if it was **Not evaluated** at time of diagnosis. If there were checked organs other than from the list, check the **Other** box and specify the organ, indicating if it is involved (select **Yes**) or not (select **No**).

See the list below:

- Skin;
- CNS;
- Testes/Ovaries;
- Other; specify.

Bibliography

1. International Agency for Research on Cancer. WHO classification of tumours of haematopoietic and lymphoid tissues: Vol. 2. 4th ed. Swerdlow SH, editor. IARC; 2017.
2. Neame PB, Soamboonsrup P, Browman GP, Meyer RM, Bengner A, Wilson WE, et al. Classifying acute leukemia by immunophenotyping: a combined FAB- immunologic classification of AML. *Blood* [Internet]. 1986;68(6):1355–62. Available from: <http://dx.doi.org/10.1182/blood.v68.6.1355.1355>