

Treatment non-HCT/CT/GT/IST therapy

**Draft guide to the completion of the EBMT data collection
form: Treatment for diagnosis_v1.0**

May 2024

EBMT Registry

.EBMT Clinical Research & Registry Department



**Co-funded by
the European Union**

Co-funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or European Health and Digital Executive Agency (HADEA). Neither the European Union nor the granting authority can be held responsible for them.

Table of Contents

Treatment non-HCT/CT/GT/IST	4
Date treatment started	4
Diagnosis for which this treatment was given	4
Line of treatment	5
Reason for this line of treatment	5
Type of relapse prevention (for malignant disease only)	5
Chemotherapy / Drug Regimen	5
Chemotherapy/Drugs	5
Start date	6
Treatment stopped	6
End date	6
Reason for treatment withdrawal	6
Chemo/Drug regimen	6
Interventions	7
Radiotherapy	7
Start date	7
Treatment stopped	7
End date	7
Splenic irradiation	7
If patient received splenic irradiation	7
Total prescribed radiation dose as per protocol	8
Number of fractions	8
Number of radiation days	8
Surgery	8
Date	8
Surgery type	8
Response To This Line Of Treatment (Disease Specific)	9
Response assessment date	9
Acute leukaemias (AML, PLN, Other)	9
Minimal residual disease (MRD)	9
Chronic leukaemias (CML, CLL, PLL, Other)	9
Chronic Myeloid Leukaemia (CML)	9
Haematological remission	10
Cytogenetic remission	10
Molecular remission	11
Chronic Lymphocytic Leukaemia (CLL), Prolymphocytic Leukaemia(PLL) and other chronic leukaemias	11
Progression sensitivity	14
Minimal residual disease (MRD)	14
Plasma cell neoplasms (PCN)	14
Number	17

Leukaemias (acute and chronic) and PCN	17
Minimal residual disease (MRD)	17
Myeloproliferative neoplasms (MPN), Myelodysplastic neoplasms (MDS), MDS/MPN overlap syndromes	17
Number	20
Lymphomas	20
Complete remission: confirmed	21
Solid tumours	21
Complete remission: confirmed	22
Relapse: resistant or sensitive	22
Other diagnosis	23

Treatment non-HCT/CT/GT/IST

The treatment for diagnosis form should be filled and submitted online in the EBMT Registry database whenever a patient with malignant disorder has received a line of treatment for the indication diagnosis. This form captures information about one line of treatment. For every new line - the new form should be filled in and submitted. One line of treatment covers all cycles of the same drugs/regimen given in the same period with the same reason, and within the same disease status of the patient. The line of treatment can include chemotherapy and/or any other intervention (radiotherapy, surgery). The form should be completed if the treatment was given for the disease that was the indication for the HCT, CT, GT or IST. The form can be registered before, between or after the main treatment (HCT, CT, GT or IST). The main reason for the line of therapy can be: induction, bridging, relapse, maintenance/preventive treatment, or consolidation.

Other EBMT DCFs will reference the current form to mark what information and at which time point of the patient's treatment history should be submitted through the non-HCT/CT/GT/IST form for either core or extended data.

Mobilisation, the preparative regimen (conditioning/lymphodepletion) and GvHD preventive treatment should be reported in the HCT/CT/GT-related forms. GvHD treatment and treatment of infectious and non-infectious complications should not be reported through the current form.

The Treatment for diagnosis form includes the following main sections:

1. Treatment;
2. Chemotherapy / Drug Regimen;
3. Interventions;
4. Response To This Line Of Treatment.

Allogeneic and Autologous HCT, Cellular therapy, Gene therapy and IST (for bone marrow failure syndromes) should **not** be reported through this form.

Date treatment started

Report the date when this line of treatment started.

Diagnosis for which this treatment was given

Enter the indication diagnosis for this line of treatment. Make sure the indication diagnosis has been registered in EBMT Registry first, using the relevant indication diagnosis form.

During online data entry the indication diagnosis must be selected from the drop-down list of all registered indication diagnoses for this patient.

Line of treatment

Report the chronological number of the line of treatment.

A line of treatment covers all cycles of the same drugs/regimen given in the same period with the same reason, and within the same disease status of the patient.

If a new line of treatment is initiated, a new form should be completed, so all separate lines are represented individually on the patient timeline.

Reason for this line of treatment

Select the main indication for this line of treatment by marking the option:

- Induction;
- Bridging;
- Relapse;
- Maintenance / preventive treatment;
- Consolidation;
- Other; specify.

If the answer option Maintenance / preventive treatment is selected, report the type of relapse prevention in the following subquestion.

Type of relapse prevention (for malignant disease only)

For malignant diseases only, specify the type of relapse prevention by selecting one of the following answer options:

- MRD negative;
- MRD positive;
- MRD unknown.

Chemotherapy / Drug Regimen

Chemotherapy/Drugs

If the patient received chemotherapy or drugs, answer **Yes** and provide details in the subsequent questions.

Answer **No**, if the patient did not receive any chemotherapy or drugs in this line of treatment.

Select **Unknown** if there is no information on whether the patients received any chemotherapy or drugs in this line of treatment.

Start date

Report the start date of the chemotherapy or drug regimen in this line of treatment. Mark as **Unknown** if the date is not available.

Treatment stopped

Report if the line of treatment was stopped and provide details in subsequent questions. If treatment has not stopped at time of reporting make sure to update this data field in the EBMT Registry whenever the treatment ends.

Mark as **Unknown** if there is no information whether treatment stopped or not.

End date

Report the end date of the chemotherapy or drug regimen in this line of treatment. If multiple drugs were given, report the end date of the last administered drug.

Mark as **Unknown** if the end date is not available.

Reason for treatment withdrawal

For Chronic Lymphocytic Leukaemia, if the treatment ended, report the reason for the treatment withdrawal by selecting one of the answer options:

- Planned withdrawal
- Toxicity
- Progression or insufficient response
- Other reason; specify.

Mark this field as **Unknown** if there is no information about the reason for the chemotherapy or drug regimen withdrawal.

Chemo/Drug regimen

Specify all the names of the chemotherapy or drugs used in this line of treatment. Please consult the LIST OF CHEMOTHERAPY DRUGS/AGENTS AND REGIMENS on the EBMT website for drugs/regimens names and synonyms.

Interventions

Radiotherapy

If the patient received radiotherapy including irradiation, answer **Yes** and provide details in the subsequent questions.

Answer **No**, if the patient did not receive any radiotherapy in this line of treatment.

Select **Unknown** if there is no information on whether the patients received or not any radiotherapy in this line of treatment.

Start date

Report the start date of the radiotherapy including irradiation in this line of treatment. Mark as **Unknown** if the date is not known.

Treatment stopped

Report if the radiotherapy including irradiation was stopped and if **Yes** provide details when. If treatment has not stopped at time of reporting make sure to update this data field in the EBMT Registry whenever the radiotherapy including irradiation ends.

Mark as **Unknown** if there is no information whether treatment stopped or not.

End date

Report the end date of the radiotherapy including irradiation in this line of treatment.

Mark as **Unknown** if the date is not available.

Splenic irradiation

For patients with Myeloproliferative neoplasms as indication for the HCT/CT, report if the patient received splenic irradiation in this line of treatment by answering **Yes** (and providing details in the subsequent questions) or **No**. Mark as **Unknown** if there is no information on whether the patient received splenic irradiation or not.

If patient received splenic irradiation

Splenic irradiation, or splenic radiation therapy, is a radiation therapy directly aimed at the spleen. It aims to reduce the spleen size in the case of splenomegaly, to relieve discomfort for patients.

Total prescribed radiation dose as per protocol

If splenic irradiation was performed, specify the total prescribed radiation dose as per protocol in Gy. Mark this field as **Unknown** if there is no information on the total prescribed radiation dose.

Number of fractions

If splenic irradiation was performed, specify the number of fractions. Mark this field as **Unknown** if there is no information on the number of fractions.

Number of radiation days

If splenic irradiation was performed, specify the number of radiation days. Mark this field as **Unknown** if there is no information on the total number of radiation days (for example, if it is ongoing).

Surgery

If the patient underwent surgery as part of this line of treatment, answer **Yes** and provide details in the subsequent questions.

Answer **No**, if the patient did not undergo surgery as part of this line of treatment.

Select **Unknown** if there is no information on whether the patients underwent surgery or not as part of this line of treatment.

Date

If the patient underwent surgery, report the surgery date. Mark as **Unknown** if the surgery date is not known.

Surgery type

If the patient underwent surgery, indicate the surgery type by selecting one of the answer options:

- Splenectomy;
- Other; specify.

Mark as **Unknown** if the surgery type is not known.

Repeat the questions for all interventions used in this line of treatment.

Response To This Line Of Treatment (Disease Specific)

Response assessment date

Report the date the best response was assessed. Mark as **Unknown** if the date of assessment is not known.

Complete only the subsequent question that is relevant for the indication diagnosis for this treatment. Note that there are different best responses for the different diseases. For all diseases the options **Unknown** and **Not evaluated** can be selected. In the EBMT Registry, the system will only show the relevant options and question(s).

Acute leukaemias (AML, PLN, Other)

Select the response from the following list, consult with criteria described in table 1:

- Complete remission (CR)
- Not in complete remission

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response	
<p>Complete remission (CR) is defined as meeting all of the following response criteria for at least four weeks:</p> <ul style="list-style-type: none"> • <5% leukemic blasts in the bone marrow • No blasts with Auer rods (applies to AML only) • No extramedullary disease (e.g., CNS, soft tissue disease) 	<p>Not in complete remission: In accordance with the defined criteria for complete remission (CR), a patient would not attain complete remission if they do not fulfil at least one of the complete remission criteria.</p>

Table 1. Acute leukaemias response.

Minimal residual disease (MRD)

Report the MRD status of acute leukaemia according to the guidelines provided [below](#).

Chronic leukaemias (CML, CLL, PLL, Other)

Chronic Myeloid Leukaemia (CML)

Select the response from the list:

- Chronic phase (CP) and type of remission (haematological, cytogenetic, molecular)
- Accelerated phase

- Blast crisis

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response		
Chronic phase (CP)	Accelerated phase (AP)	Blast crisis (BC)
<ul style="list-style-type: none"> • None of the features of accelerated phase or blast crisis 	<ul style="list-style-type: none"> • Bone marrow or peripheral blood blasts 10%-19% • Peripheral blood basophils $\geq 20\%$ • Presence of additional clonal cytogenetic abnormality in Ph+ cells (ACA)^a 	<ul style="list-style-type: none"> • Bone marrow or peripheral blood blasts $\geq 20\%$ • Extramedullary blast proliferation (myeloid sarcoma) • Presence of morphologically apparent lymphoblasts (>5%) warrants consideration of lymphoblastic crisis

Table 2. International Consensus Classification (ICC) criteria for Chronic Myeloid Leukaemias.

^aSecond Ph, trisomy 8, isochromosome 17q, trisomy 19, complex karyotype, or abnormalities of 3q26.2.

Number

If the response was chronic phase (CP), accelerated phase or blast crisis, select the number of this status.

If the response was chronic phase (CP) also indicate:

Haematological remission

If the patient was in Chronic phase (CP), report if haematological remission was achieved (answer **Yes**), or not achieved (answer **No**) according to the criteria provided in table 3. Answer **Not evaluated** if it was not evaluated or **Unknown** if it cannot be verified if it was evaluated or not.

Cytogenetic remission

If the patient was in Chronic phase (CP), report if cytogenetic remission was achieved (answer **Yes**), or not achieved (answer **No**) according to the criteria provided in table 3. Answer **Not evaluated** if it was not evaluated or **Unknown** if it cannot be verified if it was evaluated or not.

Note: A patient in cytogenetic remission must be in haematological remission but could still present a molecular relapse. This is because the cytogenetic technique has a higher resolution than haematological measurements but lower resolution than molecular methods.

Molecular remission

If the patient was in Chronic phase (CP), report if molecular remission was achieved (answer **Yes**), or not achieved (answer **No**) according to the criteria provided in table 3. Answer **Not evaluated** if it was not evaluated or **Unknown** if it cannot be verified if it was evaluated or not.

Note: A patient in molecular remission must also be in cytogenetic and haematological remission. This is because molecular techniques have a higher resolution than both haematological and cytogenetic measurements.

Disease status or best response (only CP)		
Haematological remission	Cytogenetic remission	Molecular remission
Haematological remission is defined by a patient meeting all of the following: <ul style="list-style-type: none"> ● WBC < 10 x 10⁹ /L ● Haemoglobin > 11.0 g/dL ● Platelet Count < 450 x 10⁹ /L ● Normal Differential (<1% precursor cells) ● No palpable splenomegaly ● No extramedullary disease 	Cytogenetic remission is defined by: <ul style="list-style-type: none"> ● 0% t(9;22) positive metaphases together with haematological remission ● A minimum of 20 analysable metaphases must be assessed for appropriate evaluation of a cytogenetic remission. Remission should be confirmed with repeated cytogenetic analysis within 4 to 12 weeks 	Molecular remission is defined by: <ul style="list-style-type: none"> ● Cells with the BCR::ABL1 fusion protein are not detectable, in the peripheral blood and /or the bone marrow, by an assay with a sensitivity to allow detection of one t(9;22) positive cell in 10⁵ to 10⁶ RT-PCR cells. The result should be confirmed by two consecutive tests done at least 4 weeks apart.

Table 3. Definitions of haematological, cytogenetic and molecular remission for patients in chronic phase.

Chronic Lymphocytic Leukaemia (CLL), Prolymphocytic Leukaemia(PLL) and other chronic leukaemias

Select the response from the list according to the criteria provided in table 4:

- Complete Remission (CR)
- Partial Remission (PR)
- Progression

- Stable Disease (SD)

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response		
Complete Remission (CR)	<p>All of the criteria have to be met. But:</p> <ul style="list-style-type: none"> • If a patient has all CR criteria but has persistent cytopenia, the patient can be considered as a CR as an adaptation of these guidelines. • If a patient has all criteria of a CR but bone marrow evaluation has not been performed (even with persistent cytopenia), the patient can be considered as a CR as an adaptation of these guidelines. 	
Partial Remission (PR)	<p>At least 2 of the parameters of group A and 1 parameter of group B need to improve if previously abnormal; if only 1 parameter of both groups A and B is abnormal before therapy, only 1 needs to improve.</p> <p>Clinical (palpation) evaluation only without CT-scan (or alternate imaging), according to routine practice, is accepted.</p>	
Progression	<p>At least 1 of the criteria of group A or group B has to be met.</p> <p>Sequential imaging is not warranted in CLL outside clinical trials and is not required for the EBMT Registry.</p>	Resistant to the last chemotherapy regimen
		Sensitive to the last chemotherapy regimen
Stable Disease (SD)	<p>All of the criteria have to be met. Constitutional symptoms alone do not define PD.</p>	

Table 8. Chronic lymphocytic leukaemias response.

Group	Parameter	Complete Remission (CR)	Partial Remission (PR)	Stable Disease (SD)	Progressive Disease (PD)
A	Lymph nodes	None ≥ 1.5 cm	Decrease $\geq 50\%$ (from baseline)*	Change of -49% to $+49\%$	Increase $\geq 50\%$ from baseline or from response
	Liver and/or spleen size†	Spleen size < 13 cm; liver size normal	Decrease $\geq 50\%$ (from baseline)	Change of -49% to $+49\%$	Increase $\geq 50\%$ from baseline or from response
	Constitutional symptoms	None	Any	Any	Any
	Circulating lymphocyte count	Normal	Decrease $\geq 50\%$ from baseline	Change of -49% to $+49\%$	Increase $\geq 50\%$ over baseline
B	Platelet count	$\geq 100 \times 10^9/L$	$\geq 100 \times 10^9/L$ or increase $\geq 50\%$ over baseline	Change of -49 to $+49\%$	Decrease of $\geq 50\%$ from baseline secondary to CLL
	Haemoglobin	≥ 11.0 g/dL (untransfused and without erythropoietin)	≥ 11 g/dL or increase $\geq 50\%$ over baseline	Increase < 11.0 g/dL or $< 50\%$ over baseline, or decrease < 2 g/dL	Decrease of ≥ 2 g/dL from baseline secondary to CLL
	Marrow	Normocellular, no CLL cells, no B-lymphoid nodules	Presence of CLL cells, or of B-lymphoid nodules, or not done	No change in marrow infiltrate	Increase of CLL cells by $\geq 50\%$ on successive biopsies

Table 4. Response evaluation according to 2018 iwCLL criteria.

*Sum of the products of 6 or fewer lymph nodes (as evaluated by CT scans and physical examination in clinical trials or by physical examination in general practice).

†Spleen size is considered normal if < 13 cm. There is not a firmly established international consensus of the size of a normal liver; therefore, liver size should be evaluated by imaging and manual palpation. For

the EBMT Registry, clinical (palpation) evaluation only without CT-scan (or alternate imaging), according to routine practice, is accepted.

Progression sensitivity

If the disease status or best response was progression, indicate if the progression was **resistant** to the last chemotherapy regimen the patient received, or if it was **sensitive**. If this is not known, select **unknown**.

Minimal residual disease (MRD)

Report the MRD status of chronic leukaemia according to the guidelines provided [below](#).

Plasma cell neoplasms (PCN)

Select the response from the list according to the criteria provided in table 5:

- Complete remission (CR)
- Stringent complete remission (sCR)
- Very good partial remission (VGPR)
- Partial remission (PR)
- Stable disease (no change, no response/loss of response)
- Progression
- Relapse

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response	
Complete remission (CR)	<p>Absence of detectable monoclonal immunoglobulin in serum and monoclonal light chains in the urine by immunofixation. The detection of monoclonal immunoglobulin, even at low levels which are too weak to quantitate, is not a CR.</p> <ul style="list-style-type: none"> ● <5% of plasma cells in bone marrow aspirate ● Disappearance of any soft tissue plasmacytomas. ● No increase in size or number of lytic lesions if assessed (radiographic studies are not mandatory) <p>If any of the above investigations have not been done, even if the others are indicative of a CR, the status should be registered as VGPR.</p> <p>Where CR has already been attained (bone marrow evaluation included) it may not be necessary to do a bone marrow evaluation again to confirm that the patient is still in CR (all other criteria confirming CR). Therefore, the patient is still in CR.</p>

Stringent complete remission (sCR)	<p>All of the following:</p> <ul style="list-style-type: none"> • CR as defined above • normal free light (FLC) chain ratio • Absence of clonal cells in bone marrow by immunohistochemistry or immunofluorescence
Very good partial remission (VGPR)	<p>One or more of the following:</p> <ul style="list-style-type: none"> • Serum and urine M-protein detectable by immunofixation but not on electrophoresis • $\geq 90\%$ reduction in serum M-protein plus urine M-protein level $< 0.1\text{g/}$ per 24h <p>In addition, there must be no increase in size or number of lytic lesions if assessed (radiographic studies are not mandatory)</p> <p>If any of the above investigations have not been done, even if the others are indicative of a VGPR, the status should be registered as PR.</p>
Partial remission (PR)	<p>All of the following:</p> <ul style="list-style-type: none"> • $> 50\%$ reduction in serum M-protein plus reduction in 24h urinary M-protein by $\geq 90\%$ or to $< 0.2\text{g/}$ per 24h. • A reduction of more than 50% in the size of soft tissue plasmacytomas if present at pre-treatment • No increase in size or number of lytic lesions if assessed (radiographic studies are not mandatory) <p>In the absence of measurable serum and urine M-protein, the following criteria applies:</p> <ul style="list-style-type: none"> • A decrease in the difference between involved and uninvolved free light chain (FLC) of more than 50% <p>If the FLC assay cannot be measured, the following criteria apply:</p> <ul style="list-style-type: none"> • $\geq 50\%$ reduction in plasma cells provided baseline bone marrow plasma cell percentage was $\geq 30\%$ • A reduction of more than 50% in the size of soft tissue plasmacytomas if present at pre-treatment
Stable disease (no change, no response/loss of response)	<p>Does not meet the criteria for CR, VGPR, PR or progressive disease (includes the old Minimal response (MR) criteria)</p>

<p>Progression</p>	<p>One or more of the following:</p> <ul style="list-style-type: none"> ● Increase of 25% or more in measurable monoclonal immunoglobulin in serum and urine (absolute increase must be $\geq 0.5\text{g/dL}$). This is not applicable to light chain myelomas ● Increase of 25% or more in urinary light chains (absolute increase must be $\geq 0.2\text{g/ per 24h}$) ● An increase of 25% or more in bone marrow plasma cells (absolute % must be $\geq 10\%$) ● Increase of old/appearance of new osteolytic bone lesions on x-ray ● Appearance of soft tissue plasmacytoma ● Development of hypercalcemia (corrected serum calcium $> 11.5\text{ mg/dL}$ or 2.65 mmol/L) that can be attributed solely to the plasma cell disorder <p>In the absence of measurable serum and urine M-protein, the following criterium applies:</p> <ul style="list-style-type: none"> ● An increase of 25% or more in the difference between involved and uninvolved free light chain (absolute increase must be $> 0.01\text{g/dL}$ from nadir)
<p>Relapse</p>	<p>Clinical relapse requires one or more of the following criteria:</p> <ul style="list-style-type: none"> ● Direct indicators of increasing disease and/or end organ dysfunction (CRAB features) related to the underlying clonal plasma-cell proliferative disorder. ● Development of new soft tissue plasmacytomas or bone lesions ● Increase of 50% (and at least 1 cm) in the size of existing plasmacytomas or bone lesions. ● Hypercalcemia ($> 11.5\text{ mg/dL}$) ● Decrease in haemoglobin of $> 2\text{ g/dL}$ ● Rise in serum creatinine by 2 mg/dL or more

Table 5. Plasma cell neoplasms response.

Number

Select the number of the status.

Leukaemias (acute and chronic) and PCN

Minimal residual disease (MRD)

For patients in complete remission (CR) or stringent complete remission (sCR), specify the MRD, by selection one of the answer options below:

If the patient is in haematologic CR, but has evidence of disease by more sensitive assessments including molecular, flow cytometry or cytogenetic methods, mark it as **Positive**. If the MRD assay cannot detect malignant cells mark it as **Negative**. Mark it as **Not evaluated** if MRD status evaluation was not carried out during the follow-up period. If the MRD status is not known, mark as **Unknown**.

Myeloproliferative neoplasms (MPN), Myelodysplastic neoplasms (MDS), MDS/MPN overlap syndromes

Select the response from the list according to the criteria provided in the corresponding tables 6-8:

- Complete remission (CR).
- Improvement but no CR
- Primary refractory phase (no change)
- Relapse
- Progression/worsening

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

MPN response	
Complete remission (CR)	<p>The 4 following criteria must be true:</p> <ol style="list-style-type: none"> 1. Resolution of disease-related symptoms and signs including palpable hepatosplenomegaly 2. Haemoglobin (Hb) $\geq 10\text{g/dL}$, platelet $\geq 100 \times 10^9/\text{L}$ and neutrophils $\geq 1 \times 10^9/\text{L}$ 3. $<2\%$ immature myeloid cells ($<5\%$ in splenectomized patients) 4. Normal bone marrow histology and fibrosis grade no higher than 1
Improvement but no CR	<p>Requires one criterion in absence of progression:</p> <ol style="list-style-type: none"> 1. Hb increase of 2g/dL or transfusion independence 2. Spleen reduction of 50% 3. 100% increase in platelet count and absolute platelet count of at least $50 \times 10^9/\text{L}$ 4. 100% increase in absolute neutrophil count (ANC) and an ANC of at least $0.5 \times 10^9/\text{L}$

Primary refractory phase (no change)	Treatment with intent to achieve remission was given, but no remission was achieved.
Relapse	Loss of complete remission.
Progression/Worsening	<p>Requires one of the following:</p> <ol style="list-style-type: none"> 1. Progressive splenomegaly 2. Leukaemic transformation (increase of peripheral blood blast percentage of at least 20%)

Table 6. MPN response.

MDS response	
Complete remission (CR)	<p>For patients with MDS with increased blasts: Complete remission was achieved if marrow blast count was below 5% and normalisation of peripheral blood counts was observed for at least 4 weeks.</p> <p>For patients with other types of MDS: normalisation of PB counts.</p>
Improvement but no CR	<p>1) Haematological response (in patients with cytopenia)</p> <ul style="list-style-type: none"> ● If haemoglobin < 11g/dl, erythroid response (>11 g/dl); ● If platelets <100g/l, platelet response (>100 g/l); ● If neutrophils < 1g/l, neutrophil response (>1g/l); ● If >0% peripheral blasts, response when 0% peripheral blood blasts; ● If transfusion dependant (red blood cells), independence of transfusion achieved (8 weeks without transfusions); ● If transfusion dependant (platelets), independence of transfusion achieved (8 weeks without transfusions) <p>2) Marrow blast response (in patients with increased marrow blasts): A decrease of 50% in marrow blasts, but still >5% marrow blasts.</p>
Primary refractory phase (no change)	Treatment with the intent to achieve remission was given, but no remission was achieved.
Relapse	Loss of complete remission.
Progression/ Worsening	More blasts in BM than before treatment.

Table 7. MDS disease status or best response.

MDS/MPN response	
Complete remission (CR)	Marrow blast count < 5% and a normalisation of peripheral blood counts was observed for at least 4 weeks.
Improvement but no CR	Bone marrow blasts decreased by $\geq 50\%$ after pre-treatment but still > 5%. All CR criteria were abnormal before treatment.
Primary refractory phase (no change)	Treatment with intent to achieve remission was given, but no remission was achieved.
Relapse	Loss of complete remission.
Progression/Worsening	Higher blast count in the BM and/or PB than before treatment. Worsening of cytopenias (anaemia and/or thrombocytopenia). Progression from the MD- to the MP-variant of CMML.

Table 8. MDS/MPN disease status or best response.

Number

If the disease status or best response was complete remission (CR) or relapse, select the number of this status.

Lymphomas

Select the response from the list according to the criteria provided in table 9:

- Chemorefractory relapse or progression, including primary refractory disease
- Complete remission (CR)
- Partial remission
- Stable disease (no change, no response/loss of response)
- Untreated relapse (from a previous CR) or progression (from a previous PR)

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response		
Complete remission (CR)	Complete absence of disease, no signs or symptoms of the original disease.	Confirmed
		Unconfirmed
Partial response (PR) with or without prior CR	Reduction in the disease of 50% or more	
Stable disease (no change, no response/loss of response)	Less than 50% reduction in the disease burden.	
Untreated relapse from previous CR/untreated progression from previous PR	Worsening of the disease status in patients in PR or re-appearance of the lymphoma in patients in CR, such as: recurrence of disease or systemic symptoms (B-symptoms), patient remains untreated after the relapse or progression.	
Chemorefractory relapse or progression, including primary refractory disease	Does not present any of the features of any type of remission after treatment.	

Table 9. Lymphomas response.

Complete remission: confirmed

Indicate if the complete remission was **confirmed** or **unconfirmed**. Unconfirmed means a complete response with persistent scan abnormalities of unknown significance. If it is not known if the complete remission was confirmed, select **unknown**.

Solid tumours

Select the response from the list according to the criteria provided in table 10:

- Complete remission (CR)
- First partial remission
- Partial remission (PR)
- Progressive disease
- Very good partial remission (VGPR)

- Relapse
- Stable disease (no change, no response/loss of response)

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Response		
Complete remission (CR)	The patient has achieved complete absence of disease prior to the treatment and the treatment is not part of any adjuvant therapy.	“Unconfirmed” (UCR –complete response with persistent scan abnormalities of unknown significance)
		“Confirmed” (CR with No abnormalities detected in scan)
First partial remission	The patient achieved a reduction in disease of 30% or more for the first time ever, but did not achieve complete remission.	
Partial remission (PR)	The patient achieved partial remission not for the first time.	
Progressive disease (PD)	Pay attention as it can overlap with relapse- both can be indicated. At least a 20% increase in dimension of lesions	
Very good partial remission (VGPR)		
Relapse	Reappearance of disease in patients whose last disease status was complete remission.	Sensitive: patient achieves a reduction of >50% in the disease burden after treatment for this relapse.
		Resistant: patient has not achieved a reduction of more than 50% in the disease burden after treatment for this relapse.
		Untreated: patient has not been treated for this relapse
Stable disease (no change, no response/loss of response)	<u>Target Lesions:</u> Neither sufficient shrinkage to qualify for PR nor sufficient increase to qualify for PD, taking as reference the smallest sum length diameters since the treatment started. <u>Non-Target Lesions:</u> Persistence of one or more non-target lesion(s) or/and maintenance of tumour marker level above the normal limits	

Table 10. Solid tumour response.

Complete remission: confirmed

Indicate if the complete remission was **confirmed** or **unconfirmed**. Unconfirmed means a complete response with persistent scan abnormalities of unknown significance. If it is not known if the complete remission was confirmed, select **unknown**.

Relapse: resistant or sensitive

Indicate if the relapse was **resistant** or **sensitive**.

- **Resistant:** patient has not achieved a reduction of more than 50% in the disease burden after treatment for this relapse.
- **Sensitive:** patient achieves a reduction of >30% in the disease burden after treatment for this relapse.

If it is not known if the complete remission was resistant or sensitive, select **unknown**.

Other diagnosis

If the indication diagnosis for this treatment is was Other diagnosis, specify the response by selecting one of the answer options:

- **No evidence of disease** - the patient has achieved complete absence of disease, there are no signs or symptoms of the original disease described.
- **Improved**
- **Unchanged** - Patients who have not demonstrated complete absence of disease, improvement in symptoms, or deterioration of symptoms will be classified as **Unchanged**.
- **Worse**

If the response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.