

Haematopoietic cell transplantation (HCT) annual/unscheduled follow-up

**Guide to the completion v2.7 of the EBMT
data collection form:**

HCT_FU_annual_Core_Extended_v2.6

May 2026

EBMT Registry

EBMT Clinical Research & Registry Department



**Co-funded by
the European Union**

Co-funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or European Health and Digital Executive Agency (HADEA). Neither the European Union nor the granting authority can be held responsible for them.

Table of Contents

Introduction.....	15
Haematopoietic cell transplantation (HCT) annual/unscheduled follow-up.....	15
Date of follow-up.....	15
Survival status.....	15
Main cause of death.....	15
Select treatment related cause.....	16
Infectious complication.....	16
Was an autopsy performed.....	17
Best Response.....	17
Best clinical/biological response after HCT.....	17
Date best response first observed.....	17
Graft Function.....	18
Poor graft function.....	18
Date of poor graft function.....	18
Chimaerism.....	18
Chimaerism test date.....	19
Source of cells tested.....	19
Cell types and test results.....	19
Preventive Therapies.....	19
Immunosuppression during this follow-up period.....	19
Immunosuppression stopped.....	19
End date.....	19
Was secondary letermovir prophylaxis given?.....	20
Start date secondary letermovir prophylaxis.....	20
Antimicrobial prophylaxis.....	20
Did the patient receive prophylaxis for bacterial, viral or fungal infection?.....	20
If yes, what type of prophylaxis?.....	20
Antibacterial prophylaxis.....	20
Antibiotic.....	20
Final date the antibacterial prophylaxis was discontinued.....	20
Antiviral prophylaxis.....	21
Only for allogeneic HCT: Did the patient receive CMV prophylaxis other than or in addition to letermovir during this follow-up period?.....	21
Which drugs were used?.....	21
Final date CMV prophylaxis was discontinued.....	21
Only for allogeneic HCT: Did the patient receive prophylaxis for varicella-zoster virus (VZV) or herpes simplex virus (HSV) with either acyclovir or valaciclovir during this follow-up period?.....	21
Final date VZV or HSV prophylaxis was discontinued.....	21
Only for allogeneic HCT: Did the patient receive rituximab or another anti-CD20 monoclonal drug as prophylaxis for Epstein-Barr virus post-transplant lymphoproliferative disorder (EBV-PTLD) during this follow-up period?.....	22
Did the patient receive prophylaxis for hepatitis B (HBV) during this follow-up period?.....	22

- Which drugs were used?..... 22
- Final date HBV prophylaxis was discontinued..... 22
- Antifungal prophylaxis..... 22
 - Antifungal..... 22
 - Final date the antifungal prophylaxis was discontinued.....22
 - Did the patient receive prophylaxis for Pneumocystis jirovecii pneumonia (PJP) during this follow-up period?..... 22
 - Which drugs were used?..... 23
 - Final date prophylaxis was discontinued..... 23
- Complications since the Last Follow-up - GvHD..... 23
 - Did graft versus host disease (GvHD) occur during this follow-up period?..... 23
 - Did the patient receive a systemic/immunosuppressive treatment for GvHD?..... 23
 - Date treatment started..... 24
 - Treatment stopped..... 24
 - Stop date of treatment..... 24
 - Did acute GvHD occur during this follow-up period?..... 24
 - Date of onset..... 24
 - Maximum observed organ severity score.....24
 - Overall maximum grade observed.....25
 - Steroid-refractory acute GvHD..... 26
 - Date of onset..... 26
 - aGvHD resolved?..... 27
 - Date of aGvHD resolution..... 27
 - aGvHD first line treatment..... 27
 - Did the patient receive steroids as first line treatment of aGvHD during this follow-up period?..... 27
 - Steroid details during this follow-up period..... 27
 - Were other systemic drugs/strategies used to treat aGvHD in the first line during this follow-up period?..... 27
 - Name of drug/strategy..... 27
 - How did aGvHD respond to steroids during this follow-up period?..... 28
 - Steroid sensitive (SS).....28
 - Steroid refractory (SR)..... 28
 - Steroid dependent (SD)..... 28
 - Date of onset of SD aGvHD..... 28
 - Steroid refractory/dependent aGvHD..... 28
 - Did the patient receive treatment for SR/SD aGvHD during this follow-up period?..... 28
 - Overall aGvHD grade at start of SR/SD GvHD treatment..... 28
 - Organs involved at start of SR/SD GvHD treatment.....29
 - Drugs given in this line of treatment during this follow-up period (after steroid refractoriness/dependence was established).....29
 - Organ involved during the course of treatment and response to the line of treatment during this follow-up period..... 29

Did chronic GvHD occur?.....	29
Date of onset.....	29
Maximum NIH score during this period.....	30
Date maximum NIH score.....	30
Maximum observed organ severity score.....	30
Steroid-refractory chronic GvHD.....	31
Date of onset: steroid-refractory chronic GvHD.....	31
cGvHD resolved?.....	31
Date of cGvHD resolution.....	31
Was overlap syndrome observed (features of both chronic and acute GvHD).....	31
cGvHD first line treatment.....	31
Did the patient receive steroids as first line treatment of cGvHD during this follow-up period?.	31
Steroid details during this follow-up period.....	32
Were other systemic drugs/strategies used to treat cGvHD in the first line during this follow-up period?.....	32
How did cGvHD respond to steroids during this follow-up period?.....	32
Steroid sensitive (SS).....	32
Steroid refractory (SR).....	32
Steroid dependent (SD).....	32
Steroid intolerant (SI).....	32
Date of onset of SD/SI cGvHD.....	33
Steroid refractory/dependent/intolerant cGvHD.....	33
Did the patient receive treatment for SR/SD/SI cGvHD during this follow-up period?.....	33
Overall cGvHD grade at start of SR/SD/SI treatment.....	33
Organs involved at start of SR/SD/SI treatment.....	33
Drugs given in this line of treatment during this follow-up period (after steroid refractoriness/dependence/ intolerant was established).....	33
Organ involvement during the course of treatment and response to the line of treatment during this follow-up period.....	33
Complications since the last report - Non-infectious complications.....	34
Did non-infectious complications occur during the follow-up period?.....	34
Non-infectious complication observed.....	34
Event newly developed or ongoing since previous follow-up.....	35
Maximum CTCAE grade observed.....	35
Onset date.....	36
Resolved.....	36
Stop date.....	36
Was TA-TMA treatment given during this follow-up period?.....	36
TA-TMA treatment given during this follow-up period.....	36
Other TA-TMA treatments given in this line of treatment during this follow-up period.....	37
Response to this line of TA-TMA treatment during this follow-up period.....	37
Did the patient achieve complete/partial response?.....	37

If yes, date of complete response.....	37
If no, did the patient achieve partial response?.....	37
Was SOS/VOD treatment given during this follow-up period?.....	37
SOS/VOD treatment given during this follow-up period.....	37
Other SOS/VOD treatments given in this line of treatment during this follow-up period.....	38
Response to this line of SOS/VOD treatment during this follow-up period.....	38
Did the patient achieve complete/partial response?.....	38
If yes, date of complete response.....	38
If no, did the patient achieve partial response?.....	38
Additional Late Complications.....	38
Return to work/school.....	39
Infectious complications.....	39
Did infectious complications occur during the follow-up period?.....	39
Bacterial infection.....	40
New or ongoing.....	40
Start date.....	40
Type of bacteria.....	40
Pathogen.....	41
Carbapenem (any of imipenem, meropenem, doripenem) resistance/susceptibility.....	41
KPC.....	41
OXA-48.....	41
New Delhi metallo-beta-lactamase (NDM)/VIM/IMP/Other metallo-beta-lactamase.....	41
Ceftazidime avibactam (caz-avi, avycaz)/ceftazidime avibactam+aztreonam/meropenem vaborbactam/imipenem-cilastatin-relebactam/cefiderocol resistance/susceptibility.....	42
Trimethoprim/sulfamethoxazole resistance/susceptibility.....	42
Sulbactam/cefiderocol resistance/susceptibility.....	42
Ceftolozane tazobactam (zerbaxa)/ceftazidime/cefepime/piperacillin tazobactam/ciprofloxacin or levofloxacin/amikacin resistance/susceptibility.....	42
Infection with clinical implications.....	42
Infection with clinical implications, yes:.....	42
Localisation (adapted from CTCAE terms).....	43
Were abnormalities detected upon radiological assessment?.....	43
What diagnostic technique was used?.....	43
Radiology showing new or worsening pulmonary infiltrates.....	43
What diagnostic technique was used?.....	43
Was a biopsy performed?.....	43
Date of biopsy.....	44
Was this pathogen detected in biopsy?.....	44
By what technique was the pathogen detected in biopsy?.....	44
Was bronchoalveolar lavage (BAL) performed?.....	44
Date of BAL.....	44
Was this pathogen detected in BAL?.....	44
By what technique was the pathogen detected in BAL?.....	44

Was CSF obtained?.....	44
Date of CSF.....	45
Was this pathogen detected in CSF?.....	45
By what technique was the pathogen detected in CSF?.....	45
Were typical lesions seen on the eye examination?.....	45
Was the patient transferred to the ICU due to this infection?.....	45
Contributory cause of death.....	45
Viral infection.....	45
New or ongoing.....	45
Start date.....	46
Pathogen.....	46
Infection with clinical implications.....	46
Infection with clinical implications, yes.....	46
Localisation (adapted from CTCAE terms).....	46
Were abnormalities detected upon radiological assessment?.....	47
What diagnostic technique was used?.....	47
Radiology showing new or worsening pulmonary infiltrates.....	47
What diagnostic technique was used?.....	47
Was a biopsy performed?.....	47
Date of biopsy.....	48
Was this pathogen detected in biopsy?.....	48
By what technique was the pathogen detected in biopsy?.....	48
Was bronchoalveolar lavage (BAL) performed?.....	48
Date of BAL.....	48
Was this pathogen detected in BAL?.....	48
By what technique was the pathogen detected in BAL?.....	48
Was CSF obtained?.....	48
Date of CSF.....	49
Was this pathogen detected in CSF?.....	49
By what technique was the pathogen detected in CSF?.....	49
Was an endoscopy performed?.....	49
Were gastrointestinal lesions documented?.....	49
Were typical lesions seen on the eye examination?.....	49
Was haemorrhagic cystitis diagnosed?.....	49
Was HSV resistance to antiviral drugs documented?.....	49
Was the patient transferred to the ICU due to this infection?.....	49
Contributory cause of death.....	50
Pre-emptive viral therapy.....	50
Did the patient receive pre-emptive therapy for a viral infection?.....	50
If yes, for what virus?.....	50
Specify each pre-emptive therapy modality administered for each CMV treatment course...	50
CMV pre-emptive therapy start date.....	50

Antiviral(s) used..... 50

In case of refractory CMV, was a resistance test performed?..... 50

 Which mutations were identified?..... 50

 Response to pre-emptive therapy (address response to this specific treatment modality).. 51

 Viral load at the start (+/- 3 days) of pre-emptive therapy..... 51

 Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy..... 51

 Date of discontinuation of pre-emptive therapy..... 51

Specify each pre-emptive therapy modality administered for each EBV treatment course..... 51

 EBV pre-emptive therapy start date..... 51

 Therapy used..... 51

 Response to pre-emptive therapy (address response to this specific treatment modality).. 52

 Viral load at the start (+/- 3 days) of pre-emptive therapy..... 52

 Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy..... 52

 Date of discontinuation of pre-emptive therapy..... 52

Specify each pre-emptive therapy modality administered for each BKV treatment course..... 52

 BKV pre-emptive therapy start date..... 52

 Antiviral(s) used..... 52

 Response to pre-emptive therapy (address response to this specific treatment modality).. 53

 Viral load in blood at the start (+/- 3 days) of pre-emptive therapy..... 53

 Viral load in blood at the discontinuation (+/- 3 days) of pre-emptive therapy..... 53

 Date of discontinuation of pre-emptive therapy..... 53

Specify each pre-emptive therapy modality administered for each ADV treatment course.... 53

 ADV pre-emptive therapy start date..... 53

 Antiviral(s) used..... 53

 Response to pre-emptive therapy (address response to this specific treatment modality).. 54

 Viral load in stool at the start (+/- 3 days) of pre-emptive therapy..... 54

 Viral load in blood at the start (+/- 3 days) of pre-emptive therapy..... 54

 Viral load in stool at the discontinuation (+/- 3 days) of pre-emptive therapy..... 54

 Viral load in blood at the discontinuation (+/- 3 days) of pre-emptive therapy..... 54

 Date of discontinuation of pre-emptive therapy..... 54

Specify each pre-emptive therapy modality administered for each JCV treatment course..... 54

 JCV pre-emptive therapy start date..... 55

 Antiviral(s) used..... 55

 Response to pre-emptive therapy (address response to this specific treatment modality).. 55

 Viral load at the start (+/- 3 days) of pre-emptive therapy..... 55

 Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy..... 55

 Date of discontinuation of pre-emptive therapy..... 55

Specify each pre-emptive therapy modality administered for each HHV-6 treatment course. 55

HHV-6 pre-emptive therapy start date..... 56

Antiviral(s) used..... 56

Was HHV-6 subtyping done?..... 56

 What HHV-6 subtype was identified?..... 56

Was testing for chromosomally integrated HHV-6 done?..... 56

 Was chromosomally integrated HHV-6 present?..... 56

Response to pre-emptive therapy (address response to this specific treatment modality).. 56

 Viral load at the start (+/- 3 days) of pre-emptive therapy..... 56

 Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy..... 56

 Date of discontinuation of pre-emptive therapy..... 57

Treatment of end-organ viral disease..... 57

 Did the patient receive treatment for end-organ viral disease?..... 57

 If yes, for what virus?..... 57

 CMV end-organ disease treatment start date..... 57

 Antiviral(s) used..... 57

 In case of refractory CMV, was a resistance test performed?..... 57

 Which mutation(s) were identified?..... 58

 Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)..... 58

 Date of discontinuation of end-organ disease treatment..... 58

 EBV end-organ disease treatment start date..... 58

 Therapy used..... 58

 Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)..... 58

 Date of discontinuation of end-organ disease treatment..... 59

 BKV end-organ disease treatment start date..... 59

 Antiviral(s) used..... 59

 Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)..... 59

 Date of discontinuation of end-organ disease treatment..... 59

 ADV end-organ disease treatment start date..... 60

 Antiviral(s) used..... 60

 Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)..... 60

 Date of discontinuation of end-organ disease treatment..... 60

 JCV end-organ disease treatment start date..... 60

 Therapy used..... 61

 Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)..... 61

 Date of discontinuation of end-organ disease treatment..... 61

 HHV-6 end-organ disease treatment start date..... 61

 Antiviral(s) used..... 61

Was HHV-6 subtyping done?.....	61
What HHV-6 subtype was identified?.....	62
Was testing for chromosomally integrated HHV-6 done?.....	62
Was chromosomally integrated HHV-6 present?.....	62
Response to end-organ disease treatment (address response to this/these specific modality/modalities).....	62
Date of discontinuation of end-organ disease treatment.....	62
CARV end-organ disease treatment start date.....	63
Antiviral(s) used.....	63
Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities).....	63
Date of discontinuation of end-organ disease treatment.....	63
Invasive fungal infection.....	63
New or ongoing.....	63
Start date.....	64
Type of fungus.....	64
Pathogen.....	64
Voriconazole resistance/susceptibility.....	64
Infection with clinical implications.....	64
Infection with clinical implications, yes:.....	65
Localisation (adapted from CTCAE terms).....	65
Extended dataset.....	65
Were abnormalities detected upon radiological assessment?.....	65
What diagnostic technique was used?.....	65
Radiology showing new or worsening pulmonary infiltrates.....	65
What diagnostic technique was used?.....	66
Were at least one of the following detected: Nodular lesion/Halo sign/Reverse halo sign/Cavity/Tree in bud/Ground glass/Wedge-shaped and segmental or lobal consolidation.....	66
Was PCR in blood performed?.....	66
Date of PCR in blood.....	66
Was the result confirmed by a second test?.....	66
Was a biopsy performed?.....	66
Date of biopsy.....	66
Was this pathogen detected in biopsy?.....	66
By what technique was the pathogen detected in biopsy?.....	66
Was bronchoalveolar lavage (BAL) performed?.....	67
Date of BAL.....	67
Was this pathogen detected in BAL?.....	67
By what technique was the pathogen detected in BAL?.....	67
Was CSF obtained?.....	67
Date of CSF.....	67
Was this pathogen detected in CSF?.....	67

By what technique was the pathogen detected in CSF?.....	67
Was sinus fluid sampled?.....	67
Date of sinus fluid sampling.....	67
Was this fungus detected in sinus fluid?.....	68
By what technique was the pathogen detected in sinus fluid?.....	68
Was an endoscopy performed?.....	68
Were gastrointestinal lesions documented?.....	68
Were typical lesions seen on the eye examination?.....	68
Was the patient transferred to the ICU due to this infection?.....	68
Contributory cause of death.....	68
Parasitic infection.....	69
New or ongoing.....	69
Start date.....	69
Type of parasite.....	69
Pathogen.....	69
Infection with clinical implications.....	69
Infection with clinical implications, yes.....	70
Localisation (adapted from CTCAE terms).....	70
Extended dataset.....	70
Were abnormalities detected upon radiological assessment?.....	70
What diagnostic technique was used?.....	70
Radiology showing new or worsening pulmonary infiltrates.....	70
What diagnostic technique was used?.....	71
Was a biopsy performed?.....	71
Date of biopsy.....	71
Was this pathogen detected in biopsy?.....	71
By what technique was the pathogen detected in biopsy?.....	71
Was bronchoalveolar lavage (BAL) performed?.....	71
Date of BAL.....	71
Was this pathogen detected in BAL?.....	71
By what technique was the pathogen detected in BAL?.....	71
Was CSF obtained?.....	72
Date of CSF.....	72
Was this pathogen detected in CSF?.....	72
By what technique was the pathogen detected in CSF?.....	72
Were typical lesions seen on the eye examination?.....	72
Was the patient transferred to the ICU due to this infection?.....	72
Contributory cause of death.....	72
Infection with unknown pathogen.....	72
New or ongoing.....	73
Start date.....	73
Infection with clinical implications.....	73

Infection with clinical implications, yes.....	73
Localisation (adapted from CTCAE terms).....	73
Were abnormalities detected upon radiological assessment?.....	74
What diagnostic technique was used?.....	74
Radiology showing new or worsening pulmonary infiltrates.....	74
What diagnostic technique was used?.....	74
Was the patient transferred to the ICU due to this infection?.....	74
Contributory cause of death.....	74
Vaccinations.....	75
Only if allogeneic HCT: Was the patient vaccinated with recombinant zoster vaccine (RZV; Shingrix® - GlaxoSmithKline (GSK)) during this follow-up period after the HCT treatment took place?.....	75
Date of the first RZV (Shingrix®) vaccination.....	75
Date of the last RZV (Shingrix®) vaccination.....	75
Number of doses RZV (Shingrix®) vaccine.....	75
Was the patient vaccinated against Respiratory syncytial virus (RSV) during this follow-up period after the HCT treatment took place?.....	75
What type of RSV vaccine did the patient receive?.....	75
Date of RSV vaccination.....	75
Did the patient receive monoclonal antibody against RSV during this follow-up period after the HCT treatment took place?.....	76
What type of RSV antibody did the patient receive?.....	76
Date of monoclonal antibody given.....	76
Secondary Malignancies and Autoimmune Disorders.....	76
Did secondary malignancy or autoimmune disorder occur since the last follow-up?.....	76
Was it a secondary malignancy or autoimmune disorder?.....	76
Date of diagnosis.....	76
Was this disease an indication for a subsequent HCT/CT/IST?.....	76
Additional treatment.....	76
Did the patient receive any additional disease treatment since the last follow-up?.....	76
Additional cell infusions.....	77
Did the patient receive additional cell infusions?.....	77
Is this cell infusion an allogeneic boost?.....	77
Date of the allogeneic boost.....	77
Is this cell infusion an autologous boost?.....	77
Date of the autologous boost.....	77
Did the patient receive subsequent HCT/CT?.....	78
Relapse, Progression, Recurrence of disease or Significant Worsening.....	78
MRD detectable (with any method).....	78
Date of first detectable MRD.....	78
Was there a relapse, progression, recurrence of disease or significant worsening of organ function related to the primary disease since last follow-up?.....	78
Type.....	78

Date of relapse/progression/recurrence/significant worsening.....	79
In case of relapse or progression.....	79
Type of relapse (MPN and CML only).....	79
What was the disease status? (CML only).....	79
Medullary involvement (Malignant disorders only).....	80
Extramedullary involvement (Malignant disorders only).....	80
Involvement at time of relapse (If the relapse was extramedullary or both medullary and extramedullary).....	80
Disease Status (Disease specific).....	81
Disease status at this follow-up or at time of death.....	81
Pregnancy after HCT.....	81
Has the patient become pregnant or impregnated another person since the last follow-up?.....	81
Was there an attempted pregnancy since the last follow-up?.....	81
Did the pregnancy result in a live birth?.....	82
Conception method.....	82
Appendix 1 - Disease specific best response and disease status.....	83
Acute leukaemias.....	83
Acute leukaemias disease status or best response.....	83
Chronic leukaemias.....	84
Chronic myeloid leukaemia disease status or best response.....	84
Number.....	84
Haematological remission.....	84
Cytogenetic remission.....	84
Cytogenetic detection.....	85
t(9;22) positive metaphases.....	85
t(9;22) positive cells detected by FISH.....	85
Molecular remission.....	85
BCR::ABL1 variant allele frequency (VAF).....	85
Chronic lymphocytic leukaemia (CLL), prolymphocytic leukaemia (PLL) and other chronic leukaemias disease status or best response.....	86
Progression sensitivity.....	86
Plasma cell neoplasms.....	86
Disease status or best response.....	86
Number.....	87
Immunoglobulin-related (AL) amyloidosis.....	87
Organ response.....	87
Was the patient on dialysis during this follow-up period?.....	87
Started in this follow-up period.....	88
Start date.....	88
Ongoing since previous follow-up.....	88
Did dialysis stop?.....	88
End date.....	88
Minimal residual disease.....	88

Date MRD status evaluated.....	88
Sensitivity of MRD assay.....	88
Method used.....	89
Myeloproliferative neoplasms (MPN), Myelodysplastic neoplasms (MDS), MDS/MPN overlap syndromes.....	89
Disease status or best response.....	89
Number.....	89
Autoimmune disorders.....	89
Disease status or best response.....	89
Haemoglobinopathies.....	90
Thalassemia best response.....	90
Date of last transfusion.....	90
Date of first transfusion.....	90
Thalassemia disease status.....	90
Patient requires transfusions during follow-up period.....	90
Patient requires transfusions, Yes.....	91
Date of first transfusion.....	91
Number of units.....	91
Did transfusions stop?.....	91
Date of last transfusion.....	91
Sickle cell disease best response.....	91
Date of first episode.....	91
Sickling episodes occur during follow-up period.....	92
Sickling episodes occur during follow-up period, Yes.....	92
Date of first episode.....	92
Number of SCD episodes (during follow-up).....	92
Lymphomas.....	92
Disease status or best response.....	92
Technique used for disease assessment.....	92
Solid tumours.....	93
Disease status or best response.....	93
Bone marrow failures (incl. AA).....	93
Disease status or best response.....	93
Did transfusions stop during the follow-up period?.....	93
Did the patient return to transfusion dependency afterwards?.....	94
First transfusion date.....	94
Other diagnosis.....	94
Disease status or best response.....	94
Disease status (Inborn errors only).....	94
Patient height at this follow-up.....	94
Patient weight at this follow-up.....	94
Patient is attending.....	95
Immune profiling done during this follow-up period.....	95

Test date.....	95
Cell type and test results.....	95
Disease status (Inborn errors only).....	96
Select the immunomodulatory treatments the patient received in the 3 months before the follow-up.....	96
Comorbidities during this follow up period.....	96
Was the patient admitted to ICU during this follow-up period.....	96
Cell Infusion Sheet.....	97
Chronological number of CI episode for this patient.....	97
Date of the first infusion.....	97
Number of infusions within 10 weeks.....	97
Source of cells.....	97
Type of cells.....	97
Specificity of VST product.....	98
If single-virus specific, to which virus?.....	98
If multi-virus specific, to which viruses?.....	98
Product origin.....	98
Donor source.....	98
Manufacturing method.....	98
Number of doses per infection episode.....	98
Total number of doses.....	99
Dosing interval (in days).....	99
Disease status at time of this cell infusion.....	99
Indication.....	99
Response to DLI:.....	100
MRD status.....	100
Bibliography.....	101

Introduction

Please make sure you have already checked the **Introduction to the EBMT Registry Completion Guidelines** document latest version available under *Manuals and Reference Documents* section on [EBMT website](#).

Haematopoietic cell transplantation (HCT) annual/unscheduled follow-up

The follow-up of HCT patients should be recorded using the HCT annual/unscheduled form. The form may be filled as a paper version before entering into the EBMT Registry, or can be entered online into the EBMT Registry database directly. The form should be completed at the follow-up date or at time of patient death, whichever occurs first. Annual HCT follow-up is requested according to the following schedule:

- Every year, if the patient was transplanted less than 10 years ago;
- Every 2 years, if the patient was transplanted 10–20 years ago;
- Every 5 years, if the patient was transplanted more than 20 years ago.

If there are fluctuations in the disease status during the follow-up period and the centres deem it relevant, or if the patient is discharged from the centre and/or moves to another centre, an additional report may be provided between the standard reporting schedule.

Date of follow-up

Report the date that the annual/unscheduled follow-up occurred. If the patient died, enter the date of death. If the patient was lost to follow-up, enter the date the patient was last seen alive.

Survival status

Indicate if the patient is last known to be **Alive** or **Dead** on the date of follow-up previously noted. If more than two years passed since the transplant and patient is lost to follow-up then tick the box for **Lost to follow-up**.

Main cause of death

Report only one main cause of death, even if it was considered to be a combination of various causes. If the cause of death is not known, select **Unknown**. Please select one of the following main causes of death:

- **Relapse or progression/persistent disease;**
- **Secondary malignancy;**
- **CT-related** - death caused by complications or infections after cellular therapy;
- **HCT-related** - death caused by complications or infections after transplant;
- **GT-related** - death caused by complications or infections after gene therapy;
- **IST-related** - death caused by complications or infections after immunosuppressive treatment (for patients treated with Bone Marrow Failure only).

If none of the suggested options fit, select **Other cause of death** and specify the cause of death in the textbox in English.

Select treatment related cause

In the case of treatment-related cause of death, select all the answer options that apply:

- **Graft versus host disease (GvHD);**

If the main cause of death was IST-related, graft versus host disease can not be selected.

- **Non-infectious complication;**
- **Non-infectious complication;**
- **Infectious complication.**

If none of the suggested options fit, select **Other treatment-related cause of death** and specify it in the textbox in English.

Infectious complication

In the case of an infectious complication, please specify the type of infection. In case of multiple infections with different pathogens, select all the type of infection(s) that apply:

- **Bacterial infection;**
- **Viral infection;**
- **Fungal infection;**
- **Parasitic infection;**
- **Infection with unknown pathogen.**

Please note that the category “rejection/poor graft function or failure” as contributory cause of death (previously in MedAB (auto, allo and disease-specific forms)) does not exist since the cause of death following a graft failure is generally an infection. Please note that secondary graft failure as cause of death is reportable under the list of non-infectious complications (select ‘**Fatal**’).

Extended dataset

Was an autopsy performed

Check **No**, if no autopsy has been performed. Check **Yes** if autopsy is performed. Select **Unknown** if it is unknown an autopsy was performed.

Best Response

The best response only has to be completed for **the first annual follow-up**. The disease specific options for the best response can be found in [Appendix 1](#). This section is not applicable for patients receiving HCT for Inborn errors indication diagnosis.

Best clinical/biological response after HCT

Report the patient's best response achieved after HCT but before any subsequent treatment, even if the patient got worse again afterwards. Please refer to Appendix 1 on the form to select the best response that is appropriate for the diagnosis of the patient. This includes the response observed before any subsequent treatment. If the best response after the HCT has not been evaluated, select **Not evaluated**. If the best response after the HCT is unknown, select **Unknown**.

The best response is often achieved in the first 100 days after HCT. However, for some diseases the best response to HCT may take longer and shall be reported in the first annual follow-up form (e.g. PCM). For all indication diagnoses except for inherited disorders, report the best response achieved as per the date of follow-up. In case HCT was performed for inherited disorders, this section can be left blank.

If the patient had a relapse/progression post-HCT and received therapy for the disease relapse/progression, the response to that additional therapy should not be reported in this section. The best response prior to the relapse/progression should be reported here. The response should be captured before the start of unplanned treatment of underlying disease.

Date best response first observed

Report the date the best response was first observed. The response date is the date that the sample or image was taken for assessing the response. If the patient's best response was already achieved prior to HCT (eg. HCT in CR and best response CR) the first evaluation date after the HCT should be reported. If the date is unknown, select **Unknown**.

Graft Function

Poor graft function

This question is only applicable if the follow-up is being reported for an allogeneic HCT.

Poor graft function is defined as frequent dependence on blood and/or platelet transfusions and/or growth factor support in the absence of other explanations, such as disease relapse, drugs, or infection. If poor graft function was observed - select **Yes**. If poor graft function was not observed, select **No**.

The definition for poor graft function assumes that donor chimaerism is within a desirable target level.

In contrast with graft failure, poor graft function requires some degree of allogeneic graft evidence. However, the “desirable” target level depends on the time point when it’s measured, the indication for HCT (malignant versus non-malignant), intensity of the conditioning regimen, etc.

As an example, a patient with poor haematologic function but confirmation of donor source (proof of at least mixed/partial donor chimaerism), and in the absence of other explanations (disease relapse, drug toxicity, infections, GvHD, etc) represents a case of poor graft function. However, due to the dynamic nature of chimaerism, it is advisable to reassess donor chimaerism levels to confirm the results.

Note: please do not report graft failure in this section.

- If primary graft failure occurred, please report it in the Recovery section in Absolute neutrophil count (ANC) recovery
- If secondary graft failure occurred, please report this through the Non-infectious complications section.

Date of poor graft function

Report the date when the patient started requiring frequent growth factor and/or packed RBC/platelets transfusions (at least weekly transfusions and/or growth factor support for at least 4 weeks) once other causes that could explain the poor graft function such as disease relapse, drugs, or infections have been ruled out. Mark this field as **Unknown** if the date is not known.

Chimaerism

This section is only applicable for patients that received an allogeneic HCT.

Complete this section for every chimaerism test performed within this follow-up period (after previous follow-up) until complete donor chimaerism has been achieved (>95%). If the patient has mixed chimaerism (5%-95% for either one or both myeloid and lymphoid lineages), please, complete the section

if the chimaerism results change at least 10% from the previous test. This section is optional for malignant disorders.

Chimaerism test date

Report the chimaerism test date or mark it as **Unknown** if the date is not known.

Source of cells tested

Indicate if **Peripheral blood** or **Bone marrow** was used as source of cells for the chimaerism test (select all that apply).

Cell types and test results

Select each cell type that was tested and indicate the percentage of donor cells. In case any other cell types were tested, please select **Other cell type** and indicate the type of cells in the textbox in English. If the percentage of donor cells is unknown, select **Unknown**.

Preventive Therapies

This section is only applicable for patients that received an allogeneic HCT.

Immunosuppression during this follow-up period

This question is asked to know if the GvHD prevention initiated at transplant is still ongoing at this follow up.

Select **No** if the patient was not receiving preventive (immunosuppressive) therapy for GvHD post-transplant. Select **Yes** if the patient was receiving preventive therapy post-transplant. Report as **Unknown** if it is unknown if the patient was still receiving immunosuppressive GvHD preventive treatment.

Immunosuppression stopped

If immunosuppression for GvHD prevention was stopped, please report **Yes** and provide the end date of the GvHD prevention. If the therapy is ongoing, select **No**. If it is not known if the patient is still on immunosuppressive treatment, select **Unknown**.

End date

Report the date the preventive treatment for GvHD was stopped. If the patient experiences a GvHD while receiving GvHD prevention, please report the date of onset of GvHD as the end date of GvHD prevention (as this becomes GvHD treatment rather than prevention). If the stop date was not known, select **Unknown**.

*Extended dataset***Was secondary letermovir prophylaxis given?**

Indicate whether the patient received letermovir as secondary CMV prophylaxis during the follow-up period, meaning given after pre-emptive treatment of a CMV reactivation or treatment for CMV disease regardless of whether primary letermovir prophylaxis was given during the course of the most recent HCT. Select **Unknown** if this information is unavailable.

Start date secondary letermovir prophylaxis

If **Yes**, provide the start date of the letermovir regimen. Select **Unknown** if the start date is not known.

Antimicrobial prophylaxis**Did the patient receive prophylaxis for bacterial, viral or fungal infection?**

Indicate if the patient received any type of prophylaxis by answering **Yes** or **No**.

If yes, what type of prophylaxis?

Check all types of prophylaxis the patient received.

Antibacterial prophylaxis**Antibiotic**

Check all types of antibiotics that have been administered as prophylaxis.

Indicate whether the antibacterial prophylaxis was **started during this follow-up period**, or whether it is a **Ongoing from the previous follow-up** period. If started during this follow-up period, indicate the start date or select **unknown** if you do not know the start date.

Final date the antibacterial prophylaxis was discontinued

Report the date the patient last received any type of antibacterial prophylaxis, or select **Unknown** if you do not know the final date antibacterial prophylaxis was administered, or select **Ongoing** if the patient is still receiving antibacterial prophylaxis.

Antiviral prophylaxis

Only for allogeneic HCT: Did the patient receive CMV prophylaxis other than or in addition to letermovir during this follow-up period?

Choose **Yes** if the patient received any type of CMV prophylaxis that is not letermovir. Choose **No** if the patient either received no CMV prophylaxis at all or received only letermovir. Letermovir is not included in this question as letermovir prophylaxis is a separate question in the core dataset.

Which drugs were used?

Check all types of drugs that have been administered as CMV prophylaxis. Please note that letermovir is not included here, as letermovir prophylaxis is a separate question in the core dataset. Please also do *not* include letermovir in the 'other drug' category.

Final date CMV prophylaxis was discontinued

Report the date the patient last received any type of CMV prophylaxis other than letermovir, or select **Unknown** if you do not know the final date CMV prophylaxis other than letermovir was administered, or select **Ongoing** if the patient is still receiving CMV prophylaxis.

Only for allogeneic HCT: Did the patient receive prophylaxis for varicella-zoster virus (VZV) or herpes simplex virus (HSV) with either acyclovir or valaciclovir during this follow-up period?

Indicate if either acyclovir or valaciclovir has been administered as VZV or HSV prophylaxis during this follow-up period by answering **Yes** or **No**.

Final date VZV or HSV prophylaxis was discontinued

Report the date the patient last received either acyclovir or valaciclovir as VZV or HSV prophylaxis, or select **Unknown** if you do not know the final date either acyclovir or valaciclovir was last administered as VZV or HSV prophylaxis, or select **Ongoing** if the patient is still receiving either acyclovir or valaciclovir as VZV or HSV prophylaxis.

Only for allogeneic HCT: Did the patient receive rituximab or another anti-CD20 monoclonal drug as prophylaxis for Epstein-Barr virus post-transplant lymphoproliferative disorder (EBV-PTLD) during this follow-up period?

Indicate if any anti-CD20 monoclonal drug, including rituximab, has been administered as EBV-PTLD prophylaxis during this follow-up period by answering **Yes** or **No**.

Did the patient receive prophylaxis for hepatitis B (HBV) during this follow-up period?

Indicate if any type of HBV prophylaxis has been administered during this follow-up period.

Which drugs were used?

Check all types of drugs that have been administered as HBV prophylaxis.

Final date HBV prophylaxis was discontinued

Report the date the patient last received any type of HBV prophylaxis, or select **Unknown** if you do not know the final date HBV prophylaxis was administered, or select **Ongoing** if the patient is still receiving HBV prophylaxis.

Antifungal prophylaxis

Antifungal

Check all types of antifungals that have been administered as prophylaxis except for prophylaxis against *Pneumocystis jirovecii*.

Indicate whether the antifungal prophylaxis was **Started during this follow-up period**, or whether it is a **Ongoing from the previous follow-up** period. If started during this follow-up period, indicate the start date or select **Unknown** if you do not know the start date.

Final date the antifungal prophylaxis was discontinued

Report the date the patient last received any type of antifungal prophylaxis, or select **Unknown** if you do not know the final date antifungal prophylaxis was administered, or select **Ongoing** if the patient is still receiving antifungal prophylaxis.

Did the patient receive prophylaxis for *Pneumocystis jirovecii* pneumonia (PJP) during this follow-up period?

Indicate if any type of PJP prophylaxis has been administered in this follow-up period by answering **No**, **Yes** or **Unknown**.

Which drugs were used?

Check all types of drugs that have been administered as PJP prophylaxis.

Final date prophylaxis was discontinued

Report the date the patient last received any type of PJP prophylaxis, or select **Unknown** if you do not know the final date PJP prophylaxis was administered, or select **Ongoing** if the patient is still receiving PJP prophylaxis.

Complications since the Last Follow-up - GvHD

This section should only be completed if the patient received an allogeneic HCT.

Did graft versus host disease (GvHD) occur during this follow-up period?

If **No** GvHD occurred or if this information is **Unknown**, select the appropriate answer and proceed to the next section: 'Complications since the last report - Non-infectious complications'. Select **Yes** if GvHD occurred/were ongoing/resolved in this follow up period and proceed to the next question.

Graft-versus-host disease (GvHD) refers to a clinical syndrome caused by the response of transplanted donor allogeneic cells to histocompatibility antigens expressed on tissues of the transplantation recipient. Acute GvHD (aGvHD) refers to the appearance of an allogeneic inflammatory response in exclusively three organs: the skin (inflammatory maculopapular erythematous skin rash), the liver (hyperbilirubinemia due to cholestatic jaundice), and the gastro-intestinal (GI) tract (upper and/or lower GI tract manifestations). The diagnosis must occur in the absence of manifestations of chronic GvHD (cGvHD) and should ideally be supported by positive histological findings. cGvHD is based on either the presence of specific diagnostic signs or distinctive signs accompanied by additional confirmation (e.g. biopsy or other objective diagnostic test) in at least one target organ (skin & appendages, mouth, eyes, genitalia, oesophagus, lungs and muscles & fascia). Detailed definitions are described in the 2014 NIH Consensus (1) and 2018 EBMT—NIH—CIBMTR Task Force statement on standardised terminology (2).

Did the patient receive a systemic/immunosuppressive treatment for GvHD?

Indicate if the patient received a systemic immunosuppressive treatment including ECP (Extracorporeal Photopheresis) for GvHD since the last follow-up. If the answer is **Yes**, specify if Started in this follow-up period or it is Ongoing since previous follow-up.

Date treatment started

If it started in this follow-up period, report the date the systemic immunosuppressive treatment for GvHD started or mark date **Unknown** if the date is not known.

If immunosuppressive treatment was started before the GvHD (as prevention) and continued as GvHD treatment, please indicate the date of onset of GvHD (i.e. the date that the immunosuppression was first considered as GvHD treatment, as treatment cannot technically start until onset of GvHD).

Treatment stopped

Indicate whether systemic immunosuppressive treatment for GvHD was stopped or not in this follow up period by answering **No**, **Yes** or **Unknown**.

Stop date of treatment

If treatment concluded during this follow-up, report the stop date of this treatment. Mark as **Unknown** if this date is not known.

Did acute GvHD occur during this follow-up period?

Indicate if acute graft versus host disease (aGvHD) occurred within this follow-up period (including ongoing aGvHD first reported in a previous FU) by answering **No**, **Yes** or **Unknown**. If answered Yes, specify if it Started in this follow-up period or Ongoing since previous follow-up (If aGvHD was reported in the previous FU event but was not resolved at that time).

aGvHD is a consequence of donor T-cells recognizing the patient's antigens as foreign. It usually consists of dermatitis, hepatitis, and gastroenteritis. Although it usually develops within the first 100 days, it can also appear later on.

Date of onset

If aGvHD started in this follow-up period, report the date of onset. Mark as **Unknown** if this date is not known.

Note: The date of onset only needs to be reported on the first follow-up form where the aGvHD episode first occurred.

Maximum observed organ severity score

Select for each organ listed in the table the observed severity score. If another site was also affected, answer **Yes** in **Other site affected** and specify this site in the text field in English.

The maximum grade for acute graft versus host disease (aGvHD) is defined according to the stages presented by the skin, liver, lower and upper GI tracts and can be found in table 1.

Organ	Stage	Description
Skin	0	No rash attributable to acute GVHD
	1	Skin rash < 25% body surface
	2	Skin rash 25-50% body surface
	3	Skin rash >50% body surface
	4	Generalised erythroderma (> 50% BSA) plus bullous formation and desquamation >5% of BSA
Liver	0	Total serum bilirubin < 34 µmole/L (< 2 mg/dL)
	1	Total serum bilirubin 34–50 µmole/L (2 to 3 mg/dL)
	2	Total serum bilirubin 51–102µmole/L (3.1 to 6 mg/dL)
	3	Total serum bilirubin 103–255 µmole/L (6.1 to 15 mg/dL)
	4	Total serum bilirubin >255 µmole/L (> 15 mg/dL)
Lower gut	0	Diarrhoea < 500 mL/day or <3 episodes/day for adults or diarrhoea <10 mL/kg/day or <4 episodes/day for children
	1	Diarrhoea 500–999 mL/day or 3–4 episodes/day for adults or diarrhoea 10–19.9 mL/kg/day or 4–6 episodes/day for children
	2	Diarrhoea 1000–1500mL/day or 5–7 episodes/day for adults diarrhoea 20–30 mL/kg/day or 7–10 episodes/day for children
	3	Diarrhoea >1500 mL/day or >7 episodes/day for adults or diarrhoea > 30 mL/kg/day or >10 episodes/day for children
	4	Severe abdominal pain with or without ileus or grossly bloody stools (regardless of stool volume)
Upper gut	0	No or intermittent anorexia or nausea or vomiting
	1	Persistent anorexia or nausea or vomiting

Table 1. aGvHD grading system per organ (2) <http://dx.doi.org/10.1038/s41409-018-0204-7>

Overall maximum grade observed

Select the overall maximum grade that was observed during this follow-up. If it is not known which overall maximum grade was observed, select **Unknown**. Mark it as **Not evaluated** if overall maximum grade was not carried out.

The overall grade (or the stage of skin, liver and gut) should be mentioned in the patients' file. If not clearly stated, ask the treating physician. You should report the maximum grade seen during this follow-up period as calculated from table 2 according to the MAGIC consortium.

Grade							
1	Skin stage 1 or 2	AND	Liver stage 0	AND	Upper gut stage 0	AND	Lower gut stage 0
2	Skin stage 3	AND/ OR	Liver stage 1	AND/ OR	Upper Gut stage 1	AND/ OR	Lower gut stage 1
3	Notskin stage 4	AND	Liver stage 2 or 3	AND/ OR			Lower gut stage 2 or 3
4	Skin stage 4	OR	Liver stage 4	OR			Lower gut stage 4

Table 2. Overall maximum grade for aGvHD (2) <http://dx.doi.org/10.1038/s41409-018-0204-7>

Steroid-refractory acute GvHD

Indicate if the patient experienced steroid-refractory acute GvHD or not. Steroid refractory aGvHD is defined as “failure to respond to standard steroid doses (defined as progression within 3–5 days of starting treatment or an incomplete response by 7–14 days) or recurrence after initial dose reduction (steroid dependence)” as stated in the EBMT handbook (3). Mark as **Unknown** if the steroid-refractory aGvHD was unknown.

if the patient experienced steroid-refractory acute GvHD, specify if Started in this follow-up period or it is Ongoing since previous follow-up (if the aGvHD was already reported as steroid-refractory in the previous follow-up form but was not resolved at that follow-up).

Date of onset

if Started in this follow-up period, report the date that the aGvHD was first considered steroid-refractory. Please note that by definition, this cannot be the same date as the onset of aGvHD. Mark as **Unknown** if the date of onset was unknown.

Note: The date of onset only needs to be reported on the first follow-up form where the steroid-refractory aGvHD first occurred.

*Extended dataset**aGvHD resolved?*

Please indicate whether the aGvHD was resolved or not by answering **Yes** or **No**. Mark as **Unknown** if the resolved status was unknown.

Date of aGvHD resolution

If the acute GvHD resolved, please report the date on which it was thought to have resolved completely. If the date is unknown, select **Unknown**.

aGvHD first line treatment*Did the patient receive steroids as first line treatment of aGvHD during this follow-up period?*

Please indicate here if steroids were used here for treatment of aGvHD in the first line during this follow-up period. If steroids were used in prophylaxis only, but not in treatment of aGvHD, please report 'No' here.

Steroid details during this follow-up period

Please report here the steroid used in treatment, start date in this follow-up period or mark as Ongoing since previous follow-up, initial dose (in mg/kg/day), if started in this follow up period, and stop date if applicable.

If you only have the drug in mg units, please divide by the patient weight in kg on the date of start of treatment to calculate in the correct units (mg/kg). If multiple steroids were given in the first line, please report all.

Were other systemic drugs/strategies used to treat aGvHD in the first line during this follow-up period?

Please indicate if other systemic drugs/ strategies used to treat aGVHD in the first line during this follow-up period by answering **Yes** or **No**. Mark as **Unknown** if it is unknown.

Name of drug/strategy

if other systemic drugs/ strategies (other than steroids) used to treat aGVHD in the first line, please report here all the drugs/strategies that were given in the first line treatment of aGvHD. Do not report drugs that were only given as GvHD prophylaxis.

How did aGvHD respond to steroids during this follow-up period?

Please indicate the response of aGvHD to steroids based on the following definitions below:

Steroid sensitive (SS)

If a partial or complete response is achieved in aGvHD after undergoing steroid treatment, please report 'Yes' here.

Steroid refractory (SR)

Refractory: progression in any organ within 3, 4 or 5 days of therapy onset with ≥ 2 mg/Kg/day of prednisone equivalent, or failure to improve within 5 to 7 days of treatment initiation, or incomplete response after more than 28 days of immunosuppressive treatment including steroids.

Steroid dependent (SD)

Dependent: Inability to taper prednisone under 2 mg/Kg/day after an initially successful treatment of at least 7 days or as the recurrence of aGvHD activity during steroid tapering. Specify if Started in this follow-up period or it is Ongoing since previous follow-up.

Date of onset of SD aGvHD

If steroid dependent and started in this follow-up period, indicate the date of onset of steroid dependence. Please note that by definition, this cannot be the same date as onset of aGvHD. If the date is unavailable, select **Unknown**.

Steroid refractory/dependent aGvHD

Did the patient receive treatment for SR/SD aGvHD during this follow-up period?

Indicate if the patient receives a treatment for SR/SD aGvHD during this follow-up. If **yes**, indicate whether it Started in this follow-up period or Ongoing since previous follow-up.

Overall aGvHD grade at start of SR/SD GvHD treatment

Please report the overall grade of the aGvHD on the date that SR/SD treatment started if SR/SD aGvHD treatment started in this follow-up period. This is the treatment that will be reported below (first line of SR/SD aGvHD, rather than the previously reported first line treatment of aGvHD). If treatment was started before SR/SD was established and continued as the first line treatment, please report the grade of aGvHD on the date that SR/SD was established.

Organs involved at start of SR/SD GvHD treatment

As for regular GvHD, please report all involved organ stages according to the MAGIC scale. The date of start of SR/SD aGvHD treatment applies, as above.

Drugs given in this line of treatment during this follow-up period (after steroid refractoriness/dependence was established)

Please complete the table of drugs used to treat SR/SD aGvHD. Enter drugs line by line (first line for SR/SD aGvHD, then second line and any other subsequent lines). Please note that this is not first line treatment of aGvHD, but lines of treatment after SR/SD has been established. Indicate whether it Started in this follow-up period or Ongoing since previous follow-up and stop dates, if relevant.

Organ involved during the course of treatment and response to the line of treatment during this follow-up period

Please report for each organ if it was involved or not during the course of treatment, and if yes, please also report the Best response achieved to this line of treatment and the date best response assessed. If organ specific responses are not available, please report the overall response. Responses to each line of treatment should be entered separately, starting with the first line and proceeding to subsequent lines, if applicable.

Note again, that this is the response to SR/SD aGvHD, so the response to treatment after SR/SD was established. This is not asking to report the response to first line steroid treatment of aGvHD (before SR/SD onset).

Did chronic GvHD occur?

Indicate if chronic GvHD (cGvHD) occurred or not within this follow-up period (including ongoing cGvHD first reported in a previous FU) by answering **No**, **Yes** or **Unknown**.

Date of onset

If cGvHD occurred, report if it Started in this follow-up period and the date of onset, or if it is Ongoing since previous follow-up.

If the cGvHD was already reported as steroid-refractory in the previous follow-up form but was not resolved at that follow-up, it should be registered as **Ongoing since previous follow-up**.

Note: The date of onset only needs to be reported on the first follow-up form where the cGvHD episode first occurred.

Maximum NIH score during this period

Indicate if the maximum NIH score during this period was **Mild**, **Moderate** or **Severe**. If the score is unknown, select **Unknown**. If the NIH score was not evaluated during this period, select **Not evaluated**.

The NIH scoring system was first published in 2005 and was updated in 2014 and 2022. As described in the 2014 Diagnosis and Staging Working Group report (1), eight classical organs or sites (skin, mouth, eyes, lungs, musculoskeletal system, gastrointestinal tract, genitourinary tract, and liver) are considered for calculating global score.

Elements included in the proposed global scoring include both the number of organs or sites involved and the severity score within each affected organ. Indicate the maximum NIH score during this period, as per the results of these measurements. Instructions for physicians on assessing the NIH score can be found in the EBMT handbook (1,4) or table 3.

Mild cGvHD	1 or 2 organs involved with no more than score 1 AND Lung score 0
Moderate cGvHD	3 or more organs involved with no more than score 1 OR At least 1 organ (not lung) with a score of 2 OR Lung score 1
Severe cGvHD	At least 1 organ with a score of 3 OR Lung score of 2 or 3

Table 3. Assessing the maximum NIH score (1)

In 2022, the NIH consensus (5) recognized atypical manifestations of chronic GvHD, which should be reported in the section ‘**Other site affected**’ below the list of organs involved. Atypical manifestations do not contribute to the global severity score.

Date maximum NIH score

Report the date the maximum NIH score was observed or mark it as **Unknown** if it is not known.

Maximum observed organ severity score

Select for each organ in the table the observed severity score. If another site was affected, answer **Yes** in **Other site affected** and specify this site in the text field.

Use the NIH scoring system as described in [‘Maximum NIH score during this period’](#)

Steroid-refractory chronic GvHD

Indicate if the patient experienced steroid-refractory chronic GvHD by answering **No**, **Yes** or **Unknown**.

Steroid refractory cGvHD is defined as “progression of cGvHD while on prednisone at ≥ 1 mg/kg/day for 1 to 2 weeks or stable GvHD on ≥ 0.5 mg/kg/day (or 1 mg/kg every other day) of prednisone for 1 to 2 months” (2). If answered Yes, specify if it Started in this follow-up period or Ongoing since previous follow-up.

Date of onset: steroid-refractory chronic GvHD

If Started in this follow-up period, report the date that the cGvHD was first considered steroid-refractory.

Please note that by definition, this cannot be the same date as the onset of cGvHD.

Note: The date of onset only needs to be reported on the first follow-up form where the steroid-refractory cGvHD first occurred.

Extended dataset

cGvHD resolved?

Please indicate whether the cGvHD was resolved (resolution of symptoms). If the cGvHD resolution is unknown, select **Unknown**.

Date of cGvHD resolution

If the chronic GvHD was resolved, please report the date on which this occurred. Please, select **Unknown** if the date of cGvHD resolution is unknown.

Was overlap syndrome observed (features of both chronic and acute GvHD)

Overlap syndrome is defined as the occurrence of concomitant symptoms of chronic and acute GvHD. If overlap syndrome was observed, select **Yes**. If overlap syndrome was not observed, select **No**. If the observation is unknown, select **Unknown**.

Extended dataset

cGvHD first line treatment

Did the patient receive steroids as first line treatment of cGvHD during this follow-up period?

Please indicate here if steroids were used here for treatment of cGvHD in the first line during this follow-up period. If steroids were used in prophylaxis only, but not in treatment of cGvHD, please report '**No**' here. Mark as **Unknown** if this information is not known.

Steroid details during this follow-up period

Please report here the steroid used in treatment of cGvHD, if it Started in this follow-up period and start date or if it is Ongoing since previous follow-up, initial dose (in mg/kg/day) if started in this follow up period, and stop date if applicable. If you only have the drug in mg units, please divide by the patient weight in kg on the date of start of treatment to calculate in the correct units (mg/kg). If multiple steroids were given in the first line, please report all.

Were other systemic drugs/strategies used to treat cGvHD in the first line during this follow-up period?

Please report here all the drugs/strategies that were given in the first line treatment of cGvHD in this follow-up period. Do not report drugs that were only given as GvHD prophylaxis.

How did cGvHD respond to steroids during this follow-up period?

Please indicate the response of cGvHD to steroids based on the following definitions below:

Steroid sensitive (SS)

If a partial or complete response is achieved in aGvHD after undergoing steroid treatment, please report 'Yes' here.

Steroid refractory (SR)

Refractory: progression of GvHD while on prednisone at ≥ 1 mg/Kg/day for 1-2 weeks or stable GvHD while on ≥ 0.5 mg/Kg/day (or 1 mg/Kg every other day) of prednisone for 1-2 months.

Steroid dependent (SD)

Dependent: inability to control GVHD symptoms while tapering prednisone below 0.25 mg/Kg/day (or 0.5 mg/Kg every other day) in at least two individual attempts, separated by at least 8 weeks. If answered Yes, specify it it Started in this follow-up period or Ongoing since previous follow-up.

Steroid intolerant (SI)

Intolerant: Includes avascular necrosis, severe myopathy, uncontrolled diabetes mellitus, systemic viral or fungal infections. If answered Yes, specify it it Started in this follow-up period or Ongoing since previous follow-up.

Date of onset of SD/SI cGvHD

If steroid dependent/intolerant, indicate the date of onset of steroid dependence/intolerance. (Date of onset of steroid refractory cGvHD has already been reported in the Core data field). Please note that by definition, this cannot be the same date as onset of cGvHD. If the date is not known, select **Unknown**.

Steroid refractory/dependent/intolerant cGvHD*Did the patient receive treatment for SR/SD/SI cGvHD during this follow-up period?*

Indicate if the patient receives a treatment for SR/SD cGvHD during this follow-up. If **yes**, indicate whether it Started in this follow-up period or Ongoing since previous follow-up.

Overall cGvHD grade at start of SR/SD/SI treatment

if SR/SD/SI cGvHD treatment started in this follow-up period, please report the overall grade of the cGvHD on the date that SR/SD/SI treatment started. This is the treatment that will be reported below (first line of SR/SD/SI cGvHD, rather than the previously reported first line treatment of cGvHD). If treatment was started before SR/SD/SI was established and continued as the first line treatment, please report the grade of cGvHD on the date that SR/SD/SI was established.

Organs involved at start of SR/SD/SI treatment

As for regular GvHD, please report all involved organ stages according to the NIH scoring system. The date of start of SR/SD/SI cGvHD treatment applies, as above.

Drugs given in this line of treatment during this follow-up period (after steroid refractoriness/dependence/ intolerant was established)

Please complete the table of drugs used to treat SR/SD/SI cGvHD. Enter drugs line by line (first line for SR/SD/SI cGvHD, then second line and any other subsequent lines). Please note that this is not first line treatment of aGvHD, but lines of treatment after SR/SD/SI has been established. Indicate whether it Started in this follow-up period or Ongoing since previous follow-up and stop dates, if relevant.

Organ involvement during the course of treatment and response to the line of treatment during this follow-up period

Please report if each organ was involved or not at the start of treatment, and if yes, please also report the response to this line of treatment. If organ specific responses are not available, please report the overall response. Responses to each line of treatment should be entered separately, starting with the first line and proceeding to subsequent lines, if applicable.

Note again, that this is the response to SR/SD/SI cGvHD, so the response to treatment after SR/SD/SI was established. This is not asking to report the response to first line steroid treatment of aGvHD (before SR/SD/SI onset).

Complications since the last report - Non-infectious complications

Did non-infectious complications occur during the follow-up period?

If no non-infectious complication other than GvHD occurred during the follow-up period or if the complication was grade 1 or 2, select **No** and proceed to the next section. If non-infectious complications with a CTCAE grade of at least 3 occurred or graft failure, pure red cell aplasia, posterior reversible encephalopathy syndrome or VOD of any grade occurred, select **Yes** and report in the table below. Please only report toxic events here that are above Grade 2 and not linked to GvHD and/or infections. Mark **Unknown** if this information is not available.

For adverse events not listed in the table, specify them in the **Other** text field. Consult with Appendix 3 in the paper form which non-infectious complications should not be reported even for grades 3 and 4.

For secondary graft failure: secondary graft failure is defined as a decline in hematopoietic function (may involve haemoglobin and/or platelets and/or neutrophils) necessitating blood products or growth factor support, after having met the standard definition of hematopoietic (neutrophils and platelets) recovery. Donor chimerism testing is also done to confirm the suspicion of graft failure.

Non-infectious complication observed

Specify for each adverse event listed whether it was observed or not during the follow-up being recorded. The CTCAE gradings (v5) can be found on the website of the NIH (6). Please note that if an event can be reported in more than one type of adverse event, it should be reported only once in the most precise category (eg. a cerebral thrombosis should be reported only as a cerebral thrombosis and not also as a vascular event).

If a grading is dependent on hospitalisation but the patient was an inpatient at the time of onset, the centre will make the interpretation. If the patient had been an out-patient and the severity was such that the patient would have been hospitalised, grading will be selected accordingly.

Event newly developed or ongoing since previous follow-up

Indicate if the adverse event **Newly developed** in the follow-up period (i.e. started since the last follow-up event was reported and was not present at previous follow-up) or if it was **Ongoing since the previous follow-up** (i.e. the adverse event was reported at a previous follow-up and is still present at this follow-up).

Maximum CTCAE grade observed

Select for each adverse event the maximum CTCAE grade that was observed. If the grade is unknown, select **Unknown**.

Note, for the following complications there are different grading systems to be used:

- Graft failure (fatal/non-fatal)
- Pure red cell aplasia (fatal/non-fatal)
- Posterior reversible encephalopathy syndrome (considered severe if a patient was re-hospitalised, transferred to the ICU, experiences severe mental impairment or severe cerebral edema)
- Venous-occlusive disease (VOD)/ Sinusoidal obstruction syndrome (SOS) is diagnosed based on clinical criteria. The most recently proposed EBMT criteria can be found in table 4.

Adults	Children			
Classical SOS/VOD <i>In the first 21 days after HSCT:</i> Bilirubin ≥ 2 mg/dL and two of the following criteria must be present: Painful hepatomegaly Weight gain $>5\%$ Ascites Late onset SOS/VOD <i>>21 Days after HSCT:</i> Classical VOD/SOS beyond day 21 Or Histologically proven SOS/VOD OR Two or more of the following criteria must be present: Bilirubin ≥ 2 mg/dL (or $34 \mu\text{mole/L}$) Painful hepatomegaly Weight gain $>5\%$ Ascites AND Hemodynamical or/and ultrasound evidence of SOS/VOD	No limitation for time of onset of SOS/VOD The presence of two or more of the following: <ul style="list-style-type: none"> • Unexplained consumptive and transfusion-refractory thrombocytopenia • Otherwise unexplained weight gain on three consecutive days despite the use of diuretics or a weight gain $>5\%$ above baseline value • Hepatomegaly (best if confirmed by imaging) above baseline value • Ascites (best if confirmed by imaging) above baseline value • Rising bilirubin from a baseline value on 3 consecutive days or bilirubin ≥ 2 mg/dL within 72 h 			
Severity definition				
	Mild	Moderate	Severe	Very severe
Time since first clinical symptoms of SOS/VOD	>7 Days	5–7 Days	≤ 4 Days	Any time

Bilirubin ($\mu\text{mole/L}$)	≥ 34 and < 51	≥ 51 and < 85	≥ 85 and < 136	≥ 136
Bilirubin kinetics			Doubling within 48 h	
Transaminases	$\geq 2 \times$ normal	> 2 and $\leq 5 \times$ normal	> 5 and $\leq 8 \times$ normal	$> 8 \times$ Normal
Weight increase	$< 5\%$	$\geq 5\%$ and $< 10\%$	$\geq 5\%$ and $< 10\%$	$\geq 10\%$
Renal function	$< 1.2 \times$ baseline at transplant	≥ 1.2 and $< 1.5 \times$ baseline at transplant	≥ 1.5 and $< 2 \times$ baseline at transplant	$\geq 2 \times$ baseline at transplant or others signs of MOD/MOF

Table 4. Diagnostic criteria for SOS and VOD (8)

Onset date

Report the onset date when the adverse event was first observed during this follow-up period.

Extended dataset

Resolved

If the complication was resolved in the observed period, mark as **Yes** and report the resolution date. If the complication is ongoing at time of the follow-up, select **No**. If there is no information whether the complication was resolved or not mark as **Unknown**.

Resolved is defined as three consecutive days with an absolute neutrophil count (ANC) $\geq 0.5 \times 10^9/\text{L}$, along with platelet recovery, defined as a platelet count $\geq 20 \times 10^9/\text{L}$ without platelet transfusion for at least 7 consecutive days.

Stop date

Report the date the complication was resolved or mark the date as Unknown if the date is not known. For the '**Stop date**,' report the first of the three consecutive days where the ANC reaches $\geq 0.5 \times 10^9/\text{L}$.

Was TA-TMA treatment given during this follow-up period?

Indicate if treatment was given since the last follow-up was reported by answering **No**, **Yes** or **Unknown**.

TA-TMA treatment given during this follow-up period

Please complete the table of drugs used to treat TA-TMA since the last report. Enter drugs line by line, indicate whether it Started in this follow-up period or Ongoing since previous follow-up and stop dates, if applicable. Please note that TA-TMA treatment should only be reported here if it has been given during this follow-up period

Other TA-TMA treatments given in this line of treatment during this follow-up period

Please report any of the other requested procedures or therapies that were carried out as treatment of TA-TMA here, including renal replacement therapy, mechanical ventilation or exchange plasmapheresis. Specify if it Started in this follow-up period or Ongoing since previous follow-up.

Response to this line of TA-TMA treatment during this follow-up period

Did the patient achieve complete/partial response?

Please report whether the patient achieved a complete response (CR) or not per line of treatment for TA-TMA, and if so, please provide the date of this response. CR can be defined as normal LDH, no organ manifestations, high-risk TA-TMA harmonisation criteria not fulfilled anymore.

If yes, date of complete response

If the patient achieved a complete response (CR), indicate the date of complete response or mark the date Unknown if it is not known.

If no, did the patient achieve partial response?

If the patient did not achieve a complete response, please report whether the patient achieved a partial response (PR) or not by answering **No**, **Yes** or **Unknown**, and if answered Yes, please provide the date of this response or mark the date Unknown if the date is not known. PR can be defined as LDH decreased, residual organ manifestations, high-risk TA-TMA harmonisation criteria not fulfilled anymore.

Was SOS/VOD treatment given during this follow-up period?

Indicate if treatment was given since the last follow-up was reported by answering **No**, **Yes** or **Unknown**.

SOS/VOD treatment given during this follow-up period

Please complete the table of drugs used to treat SOS/VOD since the last report. Enter drugs line by line, indicate whether it Started in this follow-up period and the start date or Ongoing since previous follow-up and stop dates, if applicable. Please note that you should only report SOS/VOD treatment if it has been given during this follow-up period.

Other SOS/VOD treatments given in this line of treatment during this follow-up period

Please report any of the other requested procedures or therapies that were carried out as treatment of SOS/VOD here, including renal replacement therapy, mechanical ventilation or extracorporeal membrane oxygenation support. Specify if it Started in this follow-up period and the start date or Ongoing since previous follow-up and stop dates, if applicable.

Response to this line of SOS/VOD treatment during this follow-up period

Did the patient achieve complete/partial response?

Please report whether the patient achieved a complete response (CR) or not per line of treatment for SOS/VOD. CR can be defined as serum bilirubin <2 mg/dL, no oxygen support, eGFR >50% from baseline before VOD and no renal replacement therapy.

If yes, date of complete response

If the patient achieved a complete response (CR), indicate the date of complete response or mark the date **Unknown** if it is not known.

If no, did the patient achieve partial response?

If the patient did not achieve a complete response, please report whether the patient achieved a partial response (PR) or not by answering **No**, **Yes** or **Unknown**, and if answered Yes, please provide the date of this response or mark the date **Unknown** if the date is not known. PR can be defined as serum bilirubin increased, but >2 mg/dL, or pulmonary dysfunction, or eGFR <=50% from baseline before SOS/VOD.

Additional Late Complications

In this section, a selection of late complications are listed. For each one, please report **Yes/No/Unknown** if the complication occurred after HCT during this follow-up period.

Please note that for many of the complications, reporting is only necessary if the complication required treatment. For these complications, if they were not treated, please do not report them. These complications include cataract diagnosis, thyroid disorder, osteoporosis, born fracture, iron overload, dyslipidemia, arterial hypertension, morbid obesity, mental health disorders, cognitive function disorder. Please also report for these complications diagnosis, if requested, and that the date that is requested is the start date of treatment and not the date of diagnosis.

Please note that only additional late complications that have not been reported already in a previous follow-up should be reported.

Return to work/school

Report if the patient has returned to work/school in this follow-up period or not by answering **No**, **Yes** or **Unknown**. If answered yes, specify the involvement: if it is parttime, fulltime or unknown and the date of return to work/school.

Infectious complications

Did infectious complications occur during the follow-up period?

Answer **Yes** if any infectious complications occurred during this follow-up period. Select **No** if infectious complications have not occurred and proceed to the next section. Mark **Unknown** if this information is not available.

Infections already reported on the previous follow-up need to be reported again if they were ongoing at the previous follow-up and thus continued into this follow-up. In this case, please update the information (e.g. clinical implications/localization/resolution) if any changes have occurred.

Infections already resolved at the previous follow-up do not need to be reported, unless a reactivation occurred.

Please note that the following infections do NOT need to be reported:

- Minor ophthalmologic bacterial infections (e.g. conjunctivitis treated topically; blepharitis treated topically; stye treated topically)
- External otitis treated topically
- Otitis media treated with oral antibiotics
- Isolated lip herpes simplex
- Bacterial tonsillitis or pharyngitis treated orally
- Laryngitis without viral identification managed at home by inhalations or without any intervention
- Upper respiratory tract infection (URTI) without viral/bacterial identification managed at home
- Bilateral cervical lymph node enlargement concurrent with URTI that resolved without specific treatment, together with the resolution of URTI
- Local superficial wound infection resolved under topical antibiotics (including impetigo)
- Minor skin bacterial infections (e.g. folliculitis; acne)
- Minor fungal skin infection (e.g. candidal intertrigo treated topically)

- Diaper rash treated with local antifungals
- Candidal balanitis treated topically
- Vaginal candidiasis treated topically or with a single oral dose
- Asymptomatic bacteriuria due to a pathogen not multi-resistant
- Single low urinary tract infection treated orally without need for hospitalisation
- Phlebitis following peripheral intravascular infusion that resolved after intravascular removal without treatment with antibiotics
- Any isolate that is considered part of the normal flora of the place (oral cavity, vagina, skin, stools) except if it carries an antimicrobial resistance that has clinical implications (induce isolation precautions or a pathogen-directed therapy)
- Positive culture without clinical implications (i.e. symptoms/signs of disease; administration of pathogen-directed therapy; isolation precautions or surveillance).

Bacterial infection

Indicate if the patient had a bacterial infection in the follow-up period after the treatment (after lymphodepleting/conditioning regimen) took place by answering **No**, **Yes** or **Unknown**. Report here only bacterial infections with microbiological documentation, otherwise they shall be reported as infection with unknown pathogen.

New or ongoing

If answered Yes in the previous question, specify if bacterial infection was newly developed or was ongoing since previous assessment (follow-up).

Start date

Report the date a first positive blood or other relevant culture or diagnostic sample was obtained. In case a diagnostic sample was obtained with a delay since the symptoms of infection started – report the date when symptoms attributable to this infection started (e.g. patient with pneumonia, urine test for legionella was sent after a few days and the test result was positive).

In case an infection was found for the first time at autopsy, use the date of death, unless you can identify the date when the symptoms attributable to this infection were reported for the first time.

If the start date was already reported on the previous follow-up form (with the **Ongoing since previous assessment** option selected), the start date does not need to be reported again.

Type of bacteria

Select the type of bacteria by marking if it is '**Gram-positive**', '**Gram-negative**' or '**Other**' (see the list in Appendix 2 of the form or available in the database).

Pathogen

Select the bacterium that caused the infection from the list in Appendix 2 of the form or available in the database. Choose the most specific option. If the pathogen cannot be found, choose the 'Gram-positive bacteria other spp', 'Gram-negative bacteria other spp' or 'Bacteria other' option and enter its name in a textbox. Always report the full name of the bacterium.

Please note that some bacteria appear several times but with the emphasis on their resistance pattern (e.g. "Pseudomonas aeruginosa (PSA) (carbapenem-susceptible)" or "Pseudomonas aeruginosa (PSA) (carbapenem-resistant)" or "Pseudomonas aeruginosa (PSA) (carbapenem susceptibility not checked)").

Common commensals (most commonly coagulase-negative Staphylococci, *Micrococcus* spp., *Bacillus* spp., *Propionibacterium* spp., *Acinetobacter lwoffii*) should be reported only if there are at least two positive blood cultures.

Extended dataset

Carbapenem (any of imipenem, meropenem, doripenem) resistance/susceptibility

Only if pathogen '*Acinetobacter baumannii*', '*Acinetobacter lwoffii*', '*Acinetobacter other*', '*Citrobacter*', '*Enterobacter*', '*Escherichia coli*', '*Klebsiella pneumoniae*', '*Klebsiella other*', '*Morganella*', '*Proteus*', '*Providencia*', '*Pseudomonas aeruginosa*', '*Pseudomonas other*', '*Raoultella*' or '*Serratia*' is selected, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to carbapenem, or indicate that the susceptibility is unknown.

KPC

For carbapenem-resistant bacteria, indicate whether the result of the KPC test was '**Negative**' or '**Positive**', or mark '**Unknown**' if this information is not available.

OXA-48

For carbapenem-resistant bacteria, indicate whether the result of the OXA-48 test was '**Negative**' or '**Positive**', or mark '**Unknown**' if this information is not available.

New Delhi metallo-beta-lactamase (NDM)/VIM/IMP/Other metallo-beta-lactamase

For carbapenem-resistant bacteria, indicate whether any of the NDM, VIM, IMP or any other metallo-beta-lactamase test was '**Negative**' or '**Positive**', or mark '**Unknown**' if this information is not available.

Ceftazidime avibactam (caz-avi, avycaz)/ceftazidime avibactam+aztreonam/meropenem vaborbactam/imipenem-cilastatin-relebactam/cefiderocol resistance/susceptibility

Only if pathogen 'Citrobacter', 'Enterobacter', 'Escherichia coli', 'Klebsiella pneumoniae', 'Klebsiella other', 'Morganella', 'Proteus', 'Providencia', 'Pseudomonas aeruginosa', 'Pseudomonas other', 'Raoultella' or 'Serratia' is selected and the pathogen is carbapenem-resistant, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to any of these antibiotics, or indicate that the susceptibility is unknown.

Trimethoprim/sulfamethoxazole resistance/susceptibility

Only if pathogen 'Stenotrophomas maltophilia' is selected, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to trimethoprim/sulfamethoxazole, or indicate that the susceptibility is unknown.

Sulbactam/cefiderocol resistance/susceptibility

Only if pathogen 'Acinetobacter baumannii', 'Acinetobacter lwoffii' or 'Acinetobacter other' is selected, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to sulbactam or cefiderocol, or indicate that the susceptibility is unknown.

Ceftolozane tazobactam (zerbaxa)/ceftazidime/cefepime/piperacillin tazobactam/ciprofloxacin or levofloxacin/amikacin resistance/susceptibility

Only if pathogen 'Pseudomonas aeruginosa' is selected, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to any of these antibiotics, or indicate that the susceptibility is unknown. Regarding Pseudomonas aeruginosa susceptibility to ciprofloxacin or levofloxacin: report '**Resistant**' if resistant to any of them and '**Susceptible**' if susceptible to both of them.

Infection with clinical implications

Indicate if the infection had clinical implications or not, or mark **Unknown** if it is not possible to identify. Infection with clinical implications is at least one of the following: symptomatic infection in the relevant organ/system, or infection that requires pathogen-directed therapy.

Infection with clinical implications, yes:

Select all clinical implications of the infection during this follow-up period that apply from suggested answer options:

- Symptoms/signs of disease;
- Administration of pathogen-directed therapy.

Localisation (adapted from CTCAE terms)

Select the localisation for the infection from the list in Appendix 3 of the form or available in the database.

In the appendix, we include both **general localisation** as part of the core dataset and **detailed localisation** as part of the extended dataset. If the more specific localisation is known, please report the **general localisation** and, if requested in the extended dataset, further specify by providing the **detailed localisation**. If only the higher-level localisation is known, report only the core dataset general localisation.

Please note that the impact of an infection can differ greatly depending on its localisation, and therefore, localisation is essential and at least 1 location involved during this follow-up period must be reported.

The localisations used in the current form are adapted from CTCAE version 5.0. Additional information on the CTCAE scoring can be found on the NIH website (6).

If the clinical information available does not specify the localisation of the infection, probably the infection was asymptomatic and will not have to be reported. Otherwise, the symptoms should guide the choice.

Extended dataset

Were abnormalities detected upon radiological assessment?

If CTCAE term 'Central nervous system infection', 'Sinusitis infective', 'Esophagus or gastric infection', 'Liver site infection', 'Lower gastrointestinal infection', 'Enteritis infective', 'Other intra-abdominal infection', 'Splenic infection' or 'Urinary tract infection' is selected, indicate whether abnormalities were detected upon radiological assessment by answering **No**, **Yes** or **Unknown**.

What diagnostic technique was used?

If abnormalities were detected upon radiological assessment, indicate by what diagnostic technique.

Radiology showing new or worsening pulmonary infiltrates

If CTCAE term 'Pneumonia' is selected, indicate whether new or worsening pulmonary infiltrates were visible upon radiology by answering **No**, **Yes** or **Unknown**.

What diagnostic technique was used?

If new or worsening pulmonary infiltrates were visible upon radiology, indicate by what diagnostic technique.

Was a biopsy performed?

If the CTCAE term was **not** 'Bacteremia', (i.e. a blood infection), indicate whether a biopsy was performed.

Date of biopsy

If a biopsy was performed, indicate the date on which the biopsy was performed. If the date is not known mark as **Unknown**.

Was this pathogen detected in biopsy?

If a biopsy was performed, indicate whether the pathogen for which you are filling out this field was detected in biopsy by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in biopsy. If more than one pathogen is detected in biopsy - report each one separately.

By what technique was the pathogen detected in biopsy?

If the pathogen for which you are filling out this field was detected in biopsy, indicate by what technique.

Was bronchoalveolar lavage (BAL) performed?

If CTCAE term 'Pneumonia' or 'Tracheobronchitis infective' is selected, indicate whether BAL was performed.

Date of BAL

If BAL was performed, indicate the date of BAL. If the date is not known mark as **Unknown**.

Was this pathogen detected in BAL?

If BAL was performed, indicate whether the pathogen for which you are filling out this field was detected in BAL by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in BAL. If more than one pathogen is detected in BAL - report each one separately.

By what technique was the pathogen detected in BAL?

If the pathogen for which you are filling out this field was detected in BAL, indicate by what technique.

Was CSF obtained?

If CTCAE term 'Central nervous system infection' is selected, indicate whether CSF was obtained by answering **No**, **Yes** or **Unknown**.

Date of CSF

If CSF was obtained, indicate the date on which CSF was obtained. If the date is not known mark as **Unknown**.

Was this pathogen detected in CSF?

If CSF was obtained, indicate whether the pathogen for which you are filling out this field was detected in CSF by answering **No**, **Yes** or **Unknown**.. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in CSF. If more than one pathogen is detected in CSF - report each one separately.

By what technique was the pathogen detected in CSF?

If the pathogen for which you are filling out this field was detected in CSF, indicate by what technique.

Were typical lesions seen on the eye examination?

If CTCAE term 'Other eye infection' is selected, indicate whether typical lesions were seen on the eye examination by answering **No**, **Yes** or **Unknown**..

Was the patient transferred to the ICU due to this infection?

Indicate whether the patient was transferred to the ICU due to this infection by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if the infection for which you are filling out this field was the reason the patient was transferred to the ICU, so not if the patient was in the ICU for any other reason.

Contributory cause of death

In case the patient is deceased, indicate if the infection contributed to death.

If there was more than one bacterial infectious complication during the follow-up period, repeat these questions for the subsequent infection. Copy and fill-in the table as many times as necessary.

Viral infection

Indicate if the patient had a viral infection in the follow-up period after the treatment (after lymphodepleting/conditioning regimen) took place. Report here only infections with microbiological documentation, otherwise they shall be reported as infection with unknown pathogen.

New or ongoing

Indicate if the patient had a **Newly developed** viral infection or if it was **Ongoing since previous assessment**.

Start date

Report the date a first positive viral test (usually PCR or antigen) was obtained. In case a diagnostic sample was obtained with a delay since the symptoms of infection started – report here the date when symptoms attributable to this infection started (e.g. patient with encephalitis, with a positive PCR in cerebrospinal fluid done 10 days after symptoms started).

In case an infection was found for the first time at autopsy, use the date of death, unless you can identify the date when the symptoms attributable to this infection were reported for the first time.

In case the start date was already reported on the previous follow-up form (with the **Ongoing since previous assessment** option selected), the start date does not need to be reported again.

Pathogen

Select the virus that caused the infection from the list in Appendix 1 of the form or available in the database. Choose the most specific option. If the pathogen cannot be found, choose the ‘Viruses other’ option and enter its name in a textbox. Always report the full name of the virus.

Infection with clinical implications

Indicate if the infection had clinical implications or not, or mark **Unknown** if it is not possible to identify. Infection with clinical implications is at least one of the following: symptomatic infection in the relevant organ/system, or infection that requires pathogen-directed therapy.

Infection with clinical implications, yes

Select all clinical implications of the infection during this follow-up period that apply:

- Symptoms/signs of disease;
- Administration of pathogen-directed therapy (including pre-emptive therapy of patients without symptoms);

Localisation (adapted from CTCAE terms)

Select the localisation for the infection from the list in Appendix 3 of the form or available in the database.

In the appendix, we include both **general localisation** as part of the core dataset and **detailed localisation** as part of the extended dataset. If the more specific localisation is known, please report the **general localisation** and, if requested in the extended dataset, further specify by providing the **detailed localisation**. If only the higher-level localisation is known, report only the core dataset general localisation.

Please note that the impact of an infection can differ greatly depending on its localisation, and therefore, localisation is essential and at least 1 location involved during this follow-up period must be reported.

The localisations used in the current form are adapted from CTCAE version 5.0. Additional information on the CTCAE scoring can be found on the NIH website (6).

If the clinical information available does not specify the localisation of the infection, probably the infection was asymptomatic and will not have to be reported. Otherwise, the symptoms should guide the choice.

A special situation exists if the virus is detected in samples from the blood (whole blood, plasma, or serum), the localization should then be reported as viremia/DNAemia by selecting from the blood infections group. This is relevant for CMV, EBV, ADV (adenovirus), BKV, JCV and HHV-6, and frequently occurs in patients without symptoms. The treatment sections below are split between pre-emptive treatment given to asymptomatic patients and the virus detected only in the blood and viral infections localized to an organ (for example lung, gastrointestinal tract or the CNS).

Extended dataset

Were abnormalities detected upon radiological assessment?

If CTCAE term 'Central nervous system infection', 'Esophagus or gastric infection', 'Liver site infection', 'Lower gastrointestinal infection', 'Enteritis infective', 'Other intra-abdominal infection' or 'Urinary tract infection' is selected, indicate whether abnormalities were detected upon radiological assessment by answering **No**, **Yes** or **Unknown**.

What diagnostic technique was used?

If abnormalities were detected upon radiological assessment, indicate by what diagnostic technique.

Radiology showing new or worsening pulmonary infiltrates

If CTCAE term 'Pneumonia' is selected, indicate whether new or worsening pulmonary infiltrates were visible upon radiology by answering **No**, **Yes** or **Unknown**.

What diagnostic technique was used?

If new or worsening pulmonary infiltrates were visible upon radiology, indicate by what diagnostic technique.

Was a biopsy performed?

If the CTCAE term was **not** 'Viremia including DNAemia' indicate whether a biopsy was performed by answering **No**, **Yes** or **Unknown**.

Date of biopsy

If a biopsy was performed, indicate the date on which the biopsy was performed. If the date is not known mark it as **Unknown**.

Was this pathogen detected in biopsy?

If a biopsy was performed, indicate whether the pathogen for which you are filling out this field was detected in biopsy by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in biopsy. If more than one pathogen is detected in biopsy - report each one separately.

By what technique was the pathogen detected in biopsy?

If the pathogen for which you are filling out this field was detected in biopsy, indicate by what technique.

Was bronchoalveolar lavage (BAL) performed?

If CTCAE term 'Pneumonia' or 'Tracheobronchitis infective' is selected, indicate whether BAL was performed by answering **No**, **Yes** or **Unknown**.

Date of BAL

If BAL was performed, indicate the date of BAL. If the date is not known mark it as **Unknown**.

Was this pathogen detected in BAL?

If BAL was performed, indicate whether the pathogen for which you are filling out this field was detected in BAL by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in BAL. If more than one pathogen is detected in BAL - report each one separately.

By what technique was the pathogen detected in BAL?

If the pathogen for which you are filling out this field was detected in BAL, indicate by what technique.

Was CSF obtained?

If CTCAE term 'Central nervous system infection' is selected, indicate whether CSF was obtained by answering **No**, **Yes** or **Unknown**.

Date of CSF

If CSF was obtained, indicate the date on which CSF was obtained. If the date is not known mark it as **Unknown**.

Was this pathogen detected in CSF?

If CSF was obtained, indicate whether the pathogen for which you are filling out this field was detected in CSF by answering **No**, **Yes** or **Unknown**. Only answer '**Yes**' if this specific pathogen was detected, so not for other pathogens potentially detected in CSF. If more than one pathogen is detected in CSF - report each one separately.

By what technique was the pathogen detected in CSF?

If the pathogen for which you are filling out this field was detected in CSF, indicate by what technique.

Was an endoscopy performed?

If CTCAE term 'Lower gastrointestinal infection' or 'esophagus or gastric infection' is selected, indicate whether an endoscopy was performed by answering **No**, **Yes** or **Unknown**.

Were gastrointestinal lesions documented?

If an endoscopy was performed, indicate whether gastrointestinal lesions were documented on this endoscopy by answering **No**, **Yes** or **Unknown**.

Were typical lesions seen on the eye examination?

If CTCAE term 'Retinitis infective' or 'Other eye infection' is selected, indicate whether typical lesions were seen on the eye examination by answering **No**, **Yes** or **Unknown**.

Was haemorrhagic cystitis diagnosed?

If CTCAE term 'Urinary tract infection' is selected, indicate whether haemorrhagic cystitis was diagnosed by answering **No**, **Yes** or **Unknown**.

Was HSV resistance to antiviral drugs documented?

If pathogen 'Herpes simplex virus (HSV)' is selected, indicate whether this virus was resistant to any antiviral drugs by answering **No**, **Yes** or **Unknown**.

Was the patient transferred to the ICU due to this infection?

Indicate whether the patient was transferred to the ICU due to this infection. Only answer '**Yes**' if the infection for which you are filling out this field was the reason the patient was transferred to the ICU, so not if the patient was in the ICU for any other reason by answering **No**, **Yes** or **Unknown**.

Contributory cause of death

In case the patient is deceased, indicate if the infection contributed to death.

Extended dataset

Pre-emptive viral therapy

Did the patient receive pre-emptive therapy for a viral infection?

Indicate whether the patient received pre-emptive therapy, meaning administration of pathogen-directed therapy in patients without symptoms.

If yes, for what virus?

If pre-emptive therapy was given, select all viruses for which pre-emptive therapy was administered.

Specify each pre-emptive therapy modality administered for each CMV treatment course

A new pre-emptive therapy course is defined either as a relapse of CMV after at least 2 weeks without antiviral therapy (success of previous treatment) or change of antiviral therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus 'CMV' is selected:

CMV pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for CMV was given. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral that was given as pre-emptive therapy for CMV, and add a new section for each antiviral. In case CMV-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

In case of refractory CMV, was a resistance test performed?

In case of refractory CMV, indicate whether a resistance test was performed by answering **No**, **Yes** or **Unknown**.

Which mutations were identified?

If a resistance test was performed, select all mutations that were identified in this resistance test.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for CMV was a **'Success'** (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a **'Failure'** (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load at the start (+/- 3 days) of pre-emptive therapy

Indicate the CMV viral load at the start of pre-emptive therapy for CMV and indicate in what unit this viral load is reported, or indicate **'Unavailable'** if the viral load is not available. If you do not know the unit, please ask your virology laboratory.

Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the CMV viral load at the discontinuation of pre-emptive therapy for CMV and indicate in what unit this viral load is reported, or indicate **'Unavailable'** if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for CMV was stopped.

If there was more than one CMV episode during the follow-up period, repeat these questions for the subsequent CMV episode. Copy and fill-in this table as many times as necessary.

Specify each pre-emptive therapy modality administered for each EBV treatment course

A new pre-emptive therapy course is defined either as a relapse of EBV after at least 2 weeks without therapy (success of previous treatment) or change of therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus 'EBV' is selected:

EBV pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for EBV was given. If the date is not known, mark as **Unknown**.

Therapy used

Select the therapy that was given as pre-emptive therapy for EBV, and add a new section for each therapy that was given. In case EBV-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for EBV was a '**Success**' (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a '**Failure**' (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load at the start (+/- 3 days) of pre-emptive therapy

Indicate the EBV viral load at the start of pre-emptive therapy for EBV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the EBV viral load at the discontinuation of pre-emptive therapy for EBV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for EBV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one EBV episode during the follow-up period, repeat these questions for the subsequent EBV episode. Copy and fill-in this table as many times as necessary.

Specify each pre-emptive therapy modality administered for each BKV treatment course

A new pre-emptive therapy course is defined either as a relapse of BKV after at least 2 weeks without antiviral therapy (success of previous treatment) or change of antiviral therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus 'BKV' is selected:

BKV pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for BKV was given. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral that was given as pre-emptive therapy for BKV, and add a new section for each antiviral that was given. In case BKV-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for BKV was a '**Success**' (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a '**Failure**' (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load in blood at the start (+/- 3 days) of pre-emptive therapy

Indicate the BKV viral load at the start of pre-emptive therapy for BKV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Viral load in blood at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the BKV viral load at the discontinuation of pre-emptive therapy for BKV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for BKV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one BKV episode during the follow-up period, repeat these questions for the subsequent BKV episode. Copy and fill-in this table as many times as necessary.

Specify each pre-emptive therapy modality administered for each ADV treatment course

A new pre-emptive therapy course is defined either as a relapse of ADV after at least 2 weeks without antiviral therapy (success of previous treatment) or change of antiviral therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus 'ADV' is selected:

ADV pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for ADV was given. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral that was given as pre-emptive therapy for ADV, and add a new section for each antiviral that was given. In case ADV-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for ADV was a ‘**Success**’ (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a ‘**Failure**’ (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load in stool at the start (+/- 3 days) of pre-emptive therapy

Indicate the ADV viral load in stool at the start of pre-emptive therapy for ADV and indicate in what unit this viral load is reported, or indicate ‘**Unavailable**’ if the viral load is not available.

Viral load in blood at the start (+/- 3 days) of pre-emptive therapy

Indicate the ADV viral load in blood at the start of pre-emptive therapy for ADV and indicate in what unit this viral load is reported, or indicate ‘**Unavailable**’ if the viral load is not available.

Viral load in stool at the discontinuation (+/- 3 days) of pre-emptive therapy

Stool can be an important sample for the detection of ADV especially in children and is used for monitoring at several centres. Indicate the ADV viral load in stool at the discontinuation of pre-emptive therapy for ADV and indicate in what unit this viral load is reported, or indicate ‘**Unavailable**’ if the viral load is not available.

Viral load in blood at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the ADV viral load in blood at the discontinuation of pre-emptive therapy for ADV and indicate in what unit this viral load is reported, or indicate ‘**Unavailable**’ if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for ADV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one ADV episode during the follow-up period, repeat these questions for the subsequent ADV episode. Copy and fill-in this table as many times as necessary.

Specify each pre-emptive therapy modality administered for each JCV treatment course

A new pre-emptive therapy course is defined either as a relapse of JCV after at least 2 weeks without antiviral therapy (success of previous treatment) or change of antiviral therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus ‘JCV’ is selected:

JCV pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for JCV was given. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral that was given as pre-emptive therapy for JCV, and add a new section for each antiviral that was given. In case JCV-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for JCV was a '**Success**' (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a '**Failure**' (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load at the start (+/- 3 days) of pre-emptive therapy

Indicate the JCV viral load at the start of pre-emptive therapy for JCV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the JCV viral load at the discontinuation of pre-emptive therapy for JCV and indicate in what unit this viral load is reported, or indicate '**Unavailable**' if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for JCV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one JCV episode during the follow-up period, repeat these questions for the subsequent JCV episode. Copy and fill-in this table as many times as necessary.

Specify each pre-emptive therapy modality administered for each HHV-6 treatment course

A new pre-emptive therapy course is defined either as a relapse of HHV-6 after at least 2 weeks without antiviral therapy (success of previous treatment) or change of antiviral therapy due to failure of any reason (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy)).

If the virus 'HHV-6' is selected:

HHV-6 pre-emptive therapy start date

Indicate the date on which the first pre-emptive therapy for HHV-6 was given. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral that was given as pre-emptive therapy for HHV-6, and add a new section for each antiviral that was given. In case HHV-6-specific T-cells (VST) were given, also fill out the additional Cell Infusion Sheet in Appendix 6.

Was HHV-6 subtyping done?

Indicate whether HHV-6 subtyping was done.

What HHV-6 subtype was identified?

If HHV-6 subtyping was done, indicate what HHV-6 subtype (A or B) was identified.

Was testing for chromosomally integrated HHV-6 done?

Indicate whether testing for chromosomally integrated HHV-6 was done.

Was chromosomally integrated HHV-6 present?

If testing for chromosomally integrated HHV-6 was done, indicate whether chromosomally integrated HHV-6 was present.

Response to pre-emptive therapy (address response to this specific treatment modality)

Indicate whether the pre-emptive therapy for HHV-6 was a **'Success'** (stopping pre-emptive therapy without need for restarting therapeutic dose of the same or another antiviral agent within 2 weeks) or a **'Failure'** (progression of viremia or stable viral load requiring change or addition of therapy, response followed by rebound (during treatment or in the 2 weeks after end of therapy), progression to disease (during treatment or in the 2 weeks after end of therapy), or unacceptable toxicity).

Viral load at the start (+/- 3 days) of pre-emptive therapy

Indicate the HHV-6 viral load at the start of pre-emptive therapy for HHV-6 and indicate in what unit this viral load is reported, or indicate **'Unavailable'** if the viral load is not available.

Viral load at the discontinuation (+/- 3 days) of pre-emptive therapy

Indicate the HHV-6 viral load at the discontinuation of pre-emptive therapy for HHV-6 and indicate in what unit this viral load is reported, or indicate **'Unavailable'** if the viral load is not available.

Date of discontinuation of pre-emptive therapy

Indicate the date on which the pre-emptive therapy for HHV-6 was stopped. If the date is not known, mark as **Unknown**.

If there was more than one HHV-6 episode during the follow-up period, repeat these questions for the subsequent HHV-6 episode. Copy and fill-in this table as many times as necessary.

Treatment of end-organ viral disease**Did the patient receive treatment for end-organ viral disease?**

Indicate whether there was administration of pathogen-directed therapy for end-organ disease caused by any viral infection.

If yes, for what virus?

If treatment for end-organ viral disease was given, select all viruses for which it was given.

Specify each treatment modality given for each occurrence of CMV disease or change in antiviral therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

It is recommended that the definitions of CMV disease according to the paper by Ljungman et al (Clin Infect Dis 2024; 79(3):787–94) are used.

If the virus 'CMV' is selected:

CMV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for CMV. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral(s) that were given as end-organ disease treatment for CMV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new antiviral. In case CMV-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

In case of refractory CMV, was a resistance test performed?

In case of refractory CMV, indicate whether a resistance test was performed.

Which mutation(s) were identified?

In case a resistance test was performed, select all mutations that were identified in this resistance test.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for CMV was a **'Success'** (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a **'Failure'** (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for CMV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one CMV episode during the follow-up period, repeat these questions for the subsequent CMV episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of EBV disease or change in therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

If the virus 'EBV' is selected:

EBV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for EBV.

Therapy used

Select the therapy/therapies that were given as end-organ disease treatment for EBV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new therapy. In case EBV-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for EBV was a **'Success'** (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a **'Failure'** (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for EBV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one EBV episode during the follow-up period, repeat these questions for the subsequent EBV episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of BKV disease or change in antiviral therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

If the virus 'BKV' is selected:

BKV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for BKV.

Antiviral(s) used

Select the antiviral(s) that were given as end-organ disease treatment for BKV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new antiviral. In case BKV-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for BKV was a '**Success**' (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a '**Failure**' (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for BKV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one BKV episode during the follow-up period, repeat these questions for the subsequent BKV episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of ADV disease or change in antiviral therapy due to failure (Defined as progression requiring change or addition of end-organ disease

treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

If the virus 'ADV' is selected:

ADV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for ADV.

Antiviral(s) used

Select the antiviral(s) that were given as end-organ disease treatment for ADV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new antiviral. In case ADV-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for ADV was a 'Success' (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a 'Failure' (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for ADV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one ADV episode during the follow-up period, repeat these questions for the subsequent ADV episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of JCV disease or change in therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

If the virus 'JCV' is selected:

JCV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for JCV. If the date is not known, mark as **Unknown**.

Therapy used

Select the therapy/therapies that were given as end-organ disease treatment for JCV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new therapy. In case JCV-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for JCV was a **'Success'** (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a **'Failure'** (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for JCV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one JCV episode during the follow-up period, repeat these questions for the subsequent JCV episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of HHV-6 disease or change in antiviral therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

If the virus 'HHV-6' is selected:

HHV-6 end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for HHV-6. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral(s) that were given as end-organ disease treatment for HHV-6. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new antiviral. In case HHV-6-specific T-cells (VST) were given, also fill in the additional Cell Infusion Sheet in Appendix 6.

Was HHV-6 subtyping done?

Indicate whether HHV-6 subtyping was done.

What HHV-6 subtype was identified?

If HHV-6 subtyping was done, indicate what HHV-6 subtype (A or B) was identified.

Was testing for chromosomally integrated HHV-6 done?

Indicate whether testing for chromosomally integrated HHV-6 was done.

Was chromosomally integrated HHV-6 present?

If testing for chromosomally integrated HHV-6 was done, indicate whether chromosomally integrated HHV-6 was present.

Response to end-organ disease treatment (address response to this/these specific modality/modalities)

Indicate whether the end-organ disease treatment for HHV-6 was a **'Success'** (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a **'Failure'** (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for HHV-6 was stopped. If the date is not known, mark as **Unknown**.

If there was more than one HHV-6 episode during the follow-up period, repeat these questions for the subsequent HHV-6 episode. Copy and fill-in this table as many times as necessary.

Specify each treatment modality given for each occurrence of community acquired respiratory virus (CARV) disease or change in antiviral therapy due to failure (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)) that occurred

As end-organ disease treatment for ADV is already asked separately, ADV is not included in the CARVs. CARVs include 'Human coronavirus (excluding SARS-CoV-2 or COVID-19)', 'Human metapneumovirus (hMPV)', 'Influenza A virus (including birdflu)', 'Influenza B virus', 'Parainfluenza', 'Respiratory syncytial virus (RSV)' and 'SARS-CoV-2 virus (COVID-19)'.

If the virus 'Community-acquired respiratory virus (CARV)' is selected:

CARV end-organ disease treatment start date

Indicate the start date of the first end-organ disease treatment given for CARV. If the date is not known, mark as **Unknown**.

Antiviral(s) used

Select the antiviral(s) that were given as end-organ disease treatment for CARV. More than one modality can be added if the intention is combination therapy and the modalities are started no more than 48 hours apart. Otherwise, add a new section for the new antiviral.

Response to end-organ disease treatment (address response to this/these specific treatment modality/modalities)

Indicate whether the end-organ disease treatment for CARV was a **'Success'** (Defined as stopping end-organ disease treatment without the need for restarting the same or other treatment) or a **'Failure'** (Defined as progression requiring change or addition of end-organ disease treatment, response followed by rebound (worsening symptoms after initial improvement, death or unacceptable toxicity)).

Date of discontinuation of end-organ disease treatment

Indicate the date on which end-organ disease treatment for CARV was stopped. If the date is not known, mark as **Unknown**.

If there was more than one CARV episode during the follow-up period, repeat these questions for the subsequent CARV episode. Copy and fill-in this table as many times as necessary.

If there was more than one viral infectious complication during the follow-up period, repeat these questions for the subsequent infection. Copy and fill-in the table as many times as necessary.

Invasive fungal infection

Indicate if the patient had a fungal infection in the follow-up period after the treatment (after lymphodepleting/conditioning regimen) took place by answering **No**, **Yes** or **Unknown**. Report here only infections with microbiological documentation, otherwise they shall be reported as infection with unknown pathogen.

New or ongoing

Indicate if the patient had a **Newly developed** fungal infection or if it was **Ongoing since previous assessment**.

Start date

Report the date a first positive culture, PCR test or galactomannan test was obtained, or the pathogen was first identified by its typical appearance in the tissue/specimen material. In case a diagnostic sample was obtained with a delay since the symptoms of infection started – report here the date when symptoms attributable to this infection started, or when this is not known, the date of the first imaging (e.g. CNS, lungs, or liver/spleen imaging for instance in hepatosplenic candidiasis in a patient with persistent fever and negative blood cultures).

In case an infection was found for the first time at autopsy, use the date of death, unless you can identify the date when the symptoms attributable to this infection were reported for the first time.

In case the start date was already reported on the previous follow-up form (with the **Ongoing since previous assessment** option selected), the start date does not need to be reported again.

Type of fungus

Select the type of fungal infection by marking if it is 'Yeasts' or 'Moulds'.

Pathogen

Select the fungus that caused the infection from the list in Appendix 2 of the form or available in the database. Choose the most specific option. If the pathogen cannot be found, choose the 'Yeasts other' or 'Moulds other' option and enter its name in a textbox. Always report the full name of the fungus. Please note that there is an option for mould infection diagnosed based on positive galactomannan only without additional microbiological confirmation.

Extended dataset

Voriconazole resistance/susceptibility

Only if pathogen 'Aspergillus flavus', 'Aspergillus fumigatus', 'Aspergillus terreus' or 'Aspergillus other' is selected, indicate whether the pathogen is '**Susceptible**' or '**Resistant**' to voriconazole, or indicate that the susceptibility is unknown.

Infection with clinical implications

Indicate if the infection had clinical implications or not, or mark unknown if it is not possible to identify. Infection with clinical implications is at least one of the following: symptomatic infection in the relevant organ/system, or infection that requires pathogen-directed therapy.

Infection with clinical implications, yes:

Select all clinical implications of the infection during this follow-up period that apply from suggested answer options:

- Symptoms/signs of disease;
- Administration of pathogen-directed therapy.

Localisation (adapted from CTCAE terms)

Select the localisation for the infection from the list in Appendix 3 of the form or available in the database.

In the appendix, we include both **general localisation** as part of the core dataset and **detailed localisation** as part of the extended dataset. If the more specific localisation is known, please report the **general localisation** and, if requested in the extended dataset, further specify by providing the **detailed localisation**. If only the higher-level localisation is known, report only the core dataset general localisation.

Please note that the impact of an infection can differ greatly depending on its localisation, and therefore, localisation is essential and at least 1 location involved during this follow-up period must be reported.

The localisations used in the current form are adapted from CTCAE version 5.0. Additional information on the CTCAE scoring can be found on the NIH website (6).

If the clinical information available does not specify the localisation of the infection, probably the infection was asymptomatic and will not have to be reported. Otherwise, the symptoms should guide the choice.

Extended dataset

Were abnormalities detected upon radiological assessment?

If CTCAE term 'Central nervous system infection', 'Sinusitis infective', 'Esophagus or gastric infection', 'Liver site infection', 'Lower gastrointestinal infection', 'Enteritis infective', 'Other intra-abdominal infection', 'Splenic infection' or 'Urinary tract infection' is selected, indicate whether abnormalities were detected upon radiological assessment.

What diagnostic technique was used?

If abnormalities were detected upon radiological assessment, indicate by what diagnostic technique.

Radiology showing new or worsening pulmonary infiltrates

If CTCAE term 'Pneumonia' is selected, indicate whether new or worsening pulmonary infiltrates were visible upon radiology.

What diagnostic technique was used?

If new or worsening pulmonary infiltrates were visible upon radiology, indicate by what diagnostic technique.

Were at least one of the following detected: Nodular lesion/Halo sign/Reverse halo sign/Cavity/Tree in bud/Ground glass/Wedge-shaped and segmental or lobal consolidation

If new or worsening pulmonary infiltrates were visible upon radiology, indicate whether any of nodular lesion, halo sign, reverse halo sign, cavity, three in bud, ground glass or wedge-shaped and segmental or global consolidation was detected.

Was PCR in blood performed?

Indicate whether PCR in blood was performed, and if so, whether the result was negative or positive.

Date of PCR in blood

If PCR in blood was performed, indicate the date on which the PCR in blood was performed.

Was the result confirmed by a second test?

If PCR in blood was performed and positive, indicate whether this positive result was confirmed by a second test.

Was a biopsy performed?

If the CTCAE term was not 'Fungemia' (i.e. a blood infection), indicate whether a biopsy was performed.

Date of biopsy

If a biopsy was performed, indicate the date on which the biopsy was performed.

Was this pathogen detected in biopsy?

If a biopsy was performed, indicate whether the pathogen for which you are filling out this field was detected in biopsy. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in biopsy. If more than one pathogen is detected in biopsy - report each one separately.

By what technique was the pathogen detected in biopsy?

If the pathogen for which you are filling out this field was detected in biopsy, indicate by what technique.

Was bronchoalveolar lavage (BAL) performed?

If CTCAE term 'Pneumonia' or 'Tracheobronchitis infective' is selected, indicate whether BAL was performed.

Date of BAL

If BAL was performed, indicate the date of BAL.

Was this pathogen detected in BAL?

If BAL was performed, indicate whether the pathogen for which you are filling out this field was detected in BAL. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in BAL. If more than one pathogen is detected in BAL - report each one separately.

By what technique was the pathogen detected in BAL?

If the pathogen for which you are filling out this field was detected in BAL, indicate by what technique.

Was CSF obtained?

If CTCAE term 'Central nervous system infection' is selected, indicate whether CSF was obtained.

Date of CSF

If CSF was obtained, indicate the date on which CSF was obtained

Was this pathogen detected in CSF?

If CSF was obtained, indicate whether the pathogen for which you are filling out this field was detected in CSF. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in CSF. If more than one pathogen is detected in CSF - report each one separately.

By what technique was the pathogen detected in CSF?

If the pathogen for which you are filling out this field was detected in CSF, indicate by what technique.

Was sinus fluid sampled?

If CTCAE term 'Sinusitis infective' is selected, indicate whether sinus fluid was sampled.

Date of sinus fluid sampling

If sinus fluid was sampled, indicate the date of sinus fluid sampling.

Was this fungus detected in sinus fluid?

If sinus fluid was sampled, indicate whether the fungus for which you are filling out this field was detected in sinus fluid. Only answer 'Yes' if this specific fungus was detected, so not for other fungi potentially detected in sinus fluid. If more than one pathogen is detected in sinus fluid - report each one separately.

By what technique was the pathogen detected in sinus fluid?

If the fungus for which you are filling out this field was detected in sinus fluid, indicate by what technique.

Was an endoscopy performed?

If CTCAE term 'Lower gastrointestinal infection' or 'esophagus or gastric infection' is selected, indicate whether an endoscopy was performed.

Were gastrointestinal lesions documented?

If an endoscopy was performed, indicate whether gastrointestinal lesions were documented on this endoscopy.

Were typical lesions seen on the eye examination?

If CTCAE term 'Retinitis infective' or 'Other eye infection' is selected, indicate whether typical lesions were seen on the eye examination.

Was the patient transferred to the ICU due to this infection?

Indicate whether the patient was transferred to the ICU due to this infection. Only answer 'Yes' if the infection for which you are filling out this field was the reason the patient was transferred to the ICU, so not if the patient was in the ICU for any other reason.

Contributory cause of death

In case the patient is deceased, indicate if the infection contributed to death.

If there was more than one fungal infectious complication during the follow-up period, repeat these questions for the subsequent infection. Copy and fill-in this table as many times as necessary.

Parasitic infection

Indicate if the patient had a parasitic infection in the follow-up period after the treatment (after lymphodepleting/conditioning regimen) took place. Report here only infections with microbiological documentation, otherwise they shall be reported as infection with unknown pathogen.

New or ongoing

Indicate if the patient had a Newly developed parasitic infection or if it was **Ongoing since previous assessment**.

Start date

Report the date a first positive antigen or DNA test was obtained or the first positive microscopic examination was performed. In case a diagnostic sample was obtained with a delay since the symptoms of infection started – report here the date when symptoms attributable to this infection started, or when this is not known, the date of the first imaging (e.g. CNS imaging for instance in Toxoplasmosis).

In case an infection was found for the first time at autopsy, use the date of death, unless you can identify the date when the symptoms attributable to this infection were reported for the first time.

In case the start date was already reported on the previous follow-up form (with the **Ongoing since previous assessment** option selected), the start date does not need to be reported again.

Type of parasite

Select the type of parasitic infection by marking if it is 'Protozoa' or 'Helminths'.

Pathogen

Select the parasite that caused the infection from the list in Appendix 2 of the form or available in the database. Choose the most specific option. If the pathogen cannot be found, choose the 'Protozoa other spp' or 'Other helminths' option and enter its name in a textbox. Always report the full name of the parasite.

Infection with clinical implications

Indicate if the infection had clinical implications or not, or mark unknown if it is not possible to identify. Infection with clinical implications is at least one of the following: symptomatic infection in the relevant organ/system, or infection that requires pathogen-directed therapy.

Infection with clinical implications, yes

Select all clinical implications of the infection during this follow-up period that apply from suggested answer options:

- Symptoms/signs of disease;
- Administration of pathogen-directed therapy.

Localisation (adapted from CTCAE terms)

Select the localisation for the infection from the list in Appendix 3 of the form or available in the database.

In the appendix, we include both **general localisation** as part of the core dataset and **detailed localisation** as part of the extended dataset. If the more specific localisation is known, please report the **general localisation** and, if requested in the extended dataset, further specify by providing the **detailed localisation**. If only the higher-level localisation is known, report only the core dataset general localisation.

Please note that the impact of an infection can differ greatly depending on its localisation, and therefore, localisation is essential and at least 1 location involved during this follow-up period must be reported.

The localisations used in the current form are adapted from CTCAE version 5.0. Additional information on the CTCAE scoring can be found on the NIH website (6).

If the clinical information available does not specify the localisation of the infection, probably the infection was asymptomatic and will not have to be reported. Otherwise, the symptoms should guide the choice.

Extended dataset

Were abnormalities detected upon radiological assessment?

If CTCAE term 'Central nervous system infection', 'Sinusitis infective', 'Esophagus or gastric infection', 'Liver site infection', 'Lower gastrointestinal infection', 'Enteritis infective', 'Other intra-abdominal infection', 'Splenic infection' or 'Urinary tract infection' is selected, indicate whether abnormalities were detected upon radiological assessment.

What diagnostic technique was used?

If abnormalities were detected upon radiological assessment, indicate by what diagnostic technique.

Radiology showing new or worsening pulmonary infiltrates

If CTCAE term 'Pneumonia' is selected, indicate whether new or worsening pulmonary infiltrates were visible upon radiology.

What diagnostic technique was used?

If new or worsening pulmonary infiltrates were visible upon radiology, indicate by what diagnostic technique.

Was a biopsy performed?

If the CTCAE term was not 'Bacteremia', 'Fungemia', 'Viremia including DNAemia' or 'DNAemia for parasitic infection' (i.e. a blood infection), indicate whether a biopsy was performed.

Date of biopsy

If a biopsy was performed, indicate the date on which the biopsy was performed.

Was this pathogen detected in biopsy?

If a biopsy was performed, indicate whether the pathogen for which you are filling out this field was detected in biopsy. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in biopsy. If more than one pathogen is detected in biopsy - report each one separately.

By what technique was the pathogen detected in biopsy?

If the pathogen for which you are filling out this field was detected in biopsy, indicate by what technique.

Was bronchoalveolar lavage (BAL) performed?

If CTCAE term 'Pneumonia' or 'Tracheobronchitis infective' is selected, indicate whether BAL was performed.

Date of BAL

If BAL was performed, indicate the date of BAL.

Was this pathogen detected in BAL?

If BAL was performed, indicate whether the pathogen for which you are filling out this field was detected in BAL. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in BAL. If more than one pathogen is detected in BAL - report each one separately.

By what technique was the pathogen detected in BAL?

If the pathogen for which you are filling out this field was detected in BAL, indicate by what technique.

Was CSF obtained?

If CTCAE term 'Central nervous system infection' is selected, indicate whether CSF was obtained.

Date of CSF

If CSF was obtained, indicate the date on which CSF was obtained

Was this pathogen detected in CSF?

If CSF was obtained, indicate whether the pathogen for which you are filling out this field was detected in CSF. Only answer 'Yes' if this specific pathogen was detected, so not for other pathogens potentially detected in CSF. If more than one pathogen is detected in CSF - report each one separately.

By what technique was the pathogen detected in CSF?

If the pathogen for which you are filling out this field was detected in CSF, indicate by what technique.

Were typical lesions seen on the eye examination?

If CTCAE term 'Retinitis infective' or 'Other eye infection' is selected, indicate whether typical lesions were seen on the eye examination.

Was the patient transferred to the ICU due to this infection?

Indicate whether the patient was transferred to the ICU due to this infection. Only answer 'Yes' if the infection for which you are filling out this field was the reason the patient was transferred to the ICU, so not if the patient was in the ICU for any other reason.

Contributory cause of death

In case the patient is deceased, indicate if the infection contributed to death.

If there was more than one parasitic infectious complication during the follow-up period, repeat these questions for the subsequent infection. Copy and fill-in this table as many times as necessary.

Infection with unknown pathogen

Indicate if the patient had an infection with unknown pathogen in the follow-up period after the treatment (after lymphodepleting/conditioning regimen) took place.

Use this section to report clinical infections without microbiological documentation, like pneumonia, cellulitis, typhlitis etc.

New or ongoing

Indicate if the patient had a **Newly developed** infection with unknown pathogen or if it was **Ongoing since previous assessment**.

Start date

Report the date the first signs or complaints were recorded or the first positive radiology was obtained. In case an infection was found for the first time at autopsy, use the date of death, unless you can identify the date when the symptoms attributable to this infection were reported for the first time.

In case the start date was already reported on the previous follow-up form (with the **Ongoing since previous assessment** option selected), the start date does not need to be reported again.

Infection with clinical implications

Infection with clinical implications is at least one of the following: symptomatic infection in the relevant organ/system, or infection that requires pathogen-directed therapy. Since an infection with an unknown pathogen always has clinical implications to be reported, the 'Infection with clinical implications, yes' field always has to be filled in.

Infection with clinical implications, yes

Select all clinical implications of the infection during this follow-up period that apply from suggested answer options:

- Symptoms/signs of disease;
- Administration of pathogen-directed therapy.

Localisation (adapted from CTCAE terms)

Select the localisation for the infection from the list in Appendix 3 of the form or available in the database.

In the appendix, we include both **general localisation** as part of the core dataset and **detailed localisation** as part of the extended dataset. If the more specific localisation is known, please report the **general localisation** and, if requested in the extended dataset, further specify by providing the **detailed localisation**. If only the higher-level localisation is known, report only the core dataset general localisation.

Please note that the impact of an infection can differ greatly depending on its localisation, and therefore, localisation is essential and at least 1 location involved during this follow-up period must be reported.

The localisations used in the current form are adapted from CTCAE version 5.0. Additional information on the CTCAE scoring can be found on the NIH website (6).

If the clinical information available doesn't specify the localisation of the infection, probably the infection was asymptomatic and will not have to be reported. Otherwise, the symptoms should guide the choice.

Extended dataset

Were abnormalities detected upon radiological assessment?

If CTCAE term 'Central nervous system infection', 'Sinusitis infective', 'Esophagus or gastric infection', 'Liver site infection', 'Lower gastrointestinal infection', 'Enteritis infective', 'Other intra-abdominal infection', 'Splenic infection' or 'Urinary tract infection' is selected, indicate whether abnormalities were detected upon radiological assessment.

What diagnostic technique was used?

If abnormalities were detected upon radiological assessment, indicate by what diagnostic technique.

Radiology showing new or worsening pulmonary infiltrates

If CTCAE term 'Pneumonia' is selected, indicate whether new or worsening pulmonary infiltrates were visible upon radiology.

What diagnostic technique was used?

If new or worsening pulmonary infiltrates were visible upon radiology, indicate by what diagnostic technique.

Was the patient transferred to the ICU due to this infection?

Indicate whether the patient was transferred to the ICU due to this infection. Only answer 'Yes' if the infection for which you are filling out this field was the reason the patient was transferred to the ICU, so not if the patient was in the ICU for any other reason.

Contributory cause of death

In case the patient is deceased, indicate if the infection contributed to death.

If there was more than one infectious complication with an unknown pathogen during the follow-up period, repeat these questions for the subsequent infection. Copy and fill-in this table as many times as necessary.

*Extended dataset***Vaccinations**

Only if allogeneic HCT: Was the patient vaccinated with recombinant zoster vaccine (RZV; Shingrix® - GlaxoSmithKline (GSK)) during this follow-up period after the HCT treatment took place?

Indicate whether the patient has been vaccinated with the RZV (Shingrix®) during this follow-up period after the HCT treatment took place.

Date of the first RZV (Shingrix®) vaccination

If the patient was vaccinated with RZV (Shingrix®), indicate on what date the patient received the first vaccination.

Date of the last RZV (Shingrix®) vaccination

If the patient was vaccinated with RZV (Shingrix®), indicate on what date the patient received the last vaccination.

Number of doses RZV (Shingrix®) vaccine

If the patient was vaccinated with RZV (Shingrix®), indicate how many doses in total the patient received.

Was the patient vaccinated against Respiratory syncytial virus (RSV) during this follow-up period after the HCT treatment took place?

Indicate whether the patient received a vaccination against the respiratory syncytial virus (RSV) during this follow-up period after the HCT treatment took place.

What type of RSV vaccine did the patient receive?

If the patient was vaccinated against RSV, indicate what type of RSV vaccine the patient received.

Date of RSV vaccination

If the patient was vaccinated against RSV, indicate on what date the patient received this vaccination.

Did the patient receive monoclonal antibody against RSV during this follow-up period after the HCT treatment took place?

Indicate whether the patient received monoclonal antibody against RSV during this follow-up period after the HCT treatment took place.

What type of RSV antibody did the patient receive?

If the patient received monoclonal antibody against RSV, indicate what type of RSV antibody the patient received.

Date of monoclonal antibody given

If the patient received monoclonal antibody against RSV, indicate on what date the patient received this monoclonal antibody.

Secondary Malignancies and Autoimmune Disorders

Did secondary malignancy or autoimmune disorder occur since the last follow-up?

Answer **No** if neither a secondary malignancy nor an autoimmune disorder has been observed after this HCT. Answer **Yes** if a secondary malignancy or an autoimmune disorder occurred and specify:

Was it a secondary malignancy or autoimmune disorder?

If answered yes in the previous question, indicate whether it is a secondary malignancy or autoimmune disorder.

Date of diagnosis

Indicate the date of diagnosis. In case the date is not known, report **Unknown**.

Was this disease an indication for a subsequent HCT/CT/IST?

If the answer is **No**, complete the non-indication diagnosis form.

If the answer is **Yes**, complete the relevant indication diagnosis form.

Additional treatment

Did the patient receive any additional disease treatment since the last follow-up?

Please indicate whether the patient received additional disease treatment (Relapse treatment, maintenance/preventive treatment, consolidation and/or other disease treatment) since the last

follow-up and specify if **Started in this follow-up period** or **Ongoing since previous follow-up**. If Started in this follow-up period, please also complete the "Treatment - non-HCT/CT/GT/IST form and /or the Cell infusion sheet".

Additional cell infusions

Did the patient receive additional cell infusions?

If the patient received additional cell infusions (i.e. MSC infusion, Donor Lymphocyte Infusion (DLI), boost etc.), excluding a new HCT and/or CT treatment, select **Yes**. If the patient did not receive additional cell infusions, select **No**. Mark **Unknown** if this information is not available.

For this question, it does not matter if it was a boost or DLI. This can be clarified in the next questions.

If answered Yes, the following should be taken into account:

If the cells were infused with the aim of improving chimaerism, or preventing or treating relapses, it most likely was a DLI. The treating physician knows the aim of the infusion. In rare cases, the aim can be as for a boost and a DLI. In this case boost should be selected and the cell infusion sheet for DLI should be completed.

Is this cell infusion an allogeneic boost?

Indicate whether the cell infusion was an allogeneic boost or not.

An allogeneic boost is an infusion of cells from the same donor without conditioning, with no evidence of graft rejection. A boost is infused with the aim of providing enough hematopoietic cells to have an effect on engraftment.

If cells are not from the same donor OR there is conditioning (chemotherapy and/or TBI), then it is considered to be an HCT and not a boost.

Date of the allogeneic boost

If applicable, report the date the boost took place.

Is this cell infusion an autologous boost?

If the cell infusion was an autologous boost answer **Yes**. If it was not an autologous boost, select **No**.

An autologous boost is an infusion of pre-collected and stored autologous stem cells without conditioning.

Date of the autologous boost

If applicable, report the date the boost took place.

Note: If this cell infusion is not a boost, attach the Cell Infusion (CI) sheet available in Appendix 6, completing as many sheets as episodes of cell infusion that took place during this follow-up period; then continue with questions below.

Did the patient receive subsequent HCT/CT?

Report whether or not the patient received a subsequent HCT/CT, either at your or another centre. Please make sure that this subsequent treatment is registered using either the HCT or CT day 0 form and the disease status HCT/CT/GT/IST day 0 form.

Relapse, Progression, Recurrence of disease or Significant Worsening

MRD detectable (with any method)

Only for Acute leukaemia, report if MRD detectable by answering No, Yes or Unknown.

Date of first detectable MRD

Report the date of the first detectable MRD or mark the date Unknown if it is not known.

Was there a relapse, progression, recurrence of disease or significant worsening of organ function related to the primary disease since last follow-up?

Indicate if there was a relapse, progression, recurrence or significant worsening of organ function related to the primary disease since the last follow-up detected by any method. If multiple instances of relapse, progression, recurrence of disease or significant worsening took place in this follow-up period, report all instances. When filling this question in the EBMT registry, click the **add** button as many times as necessary for reporting all instances. ('+ Add' on the EBMT Registry.) If the answer is **No**, proceed to the next section.

Type

Report if a **relapse or recurrence of disease** is reported, or a **(continuous) progression or significant worsening**. Relapse or Recurrence of Disease is defined as active disease after complete remission for malignant diseases, or after absence of symptoms for non-malignant diseases. (Continuous) progression or worsening is defined as active disease without complete remission since diagnosis or since last main treatment for malignant diseases, or continuous active disease without absence of symptom since last main treatment for non-malignant diseases

Date of relapse/progression/recurrence/significant worsening

Report the date of the relapse/progression/recurrence/significant worsening since HCT. If the date is not known, select **unknown**.

Extended dataset

In case of relapse or progression

Type of relapse (MPN and CML only)

Indicate if there was a relapse related to the primary disease after HCT.

Select from the list the worst disease status detected at this time point:

- Haematological;
- Cytogenetic;
- Molecular.

If the disease status at relapse is not known, select **Unknown**.

What was the disease status? (CML only)

If haematological relapse occurred, also indicate the disease status:

- Chronic phase;
- Accelerated phase;
- Blast crisis.

Disease status		
Chronic phase (CP)	Accelerated phase (AP)	Blast crisis (BC)
<ul style="list-style-type: none"> ● None of the features of accelerated phase or blast crisis 	<ul style="list-style-type: none"> ● Bone marrow or peripheral blood blasts 10%-19% ● Peripheral blood basophils $\geq 20\%$ ● Presence of additional clonal cytogenetic abnormality in Ph+ cells (ACA)^a 	<ul style="list-style-type: none"> ● Bone marrow or peripheral blood blasts $\geq 20\%$ ● Extramedullary blast proliferation (myeloid sarcoma) ● Presence of morphologically apparent lymphoblasts (>5%) warrants consideration of lymphoblastic crisis

Table 5. International Consensus Classification (ICC) criteria for Chronic Myeloid Leukaemias.

Medullary involvement (Malignant disorders only)

Indicate if the marrow or blood were affected by the disease. Please be aware that although the vast majority of acute leukaemias involve the invasion of the bone marrow by blasts, there are cases where blast invasion is only found in organs other than the bone marrow (e.g. choromas).

Medullary relapse is when malignant cells are only found in the bone marrow. In case of extramedullary relapse, malignant cells are found in sites other than the bone marrow, such as soft tissues or organs.

Extramedullary involvement (Malignant disorders only)

Extramedullary involvement (EMI) refers to disease cells found in organs or tissue outside the blood or bone marrow. The most common sites of extramedullary disease are the central nervous system (CNS), skin and ovaries/testes.

Involvement at time of relapse (If the relapse was extramedullary or both medullary and extramedullary)

Report if **Skin**, **CNS** (central nervous system) or **Testes/Ovaries** were involved at time of relapse. Also indicate if any other site was involved. If yes, please specify this in the textfield in English.

Disease Status (Disease specific)

Disease status at this follow-up or at time of death

Indicate the disease status at this follow-up or at time of death corresponding to indication diagnosis by selecting from the list provided in Appendix 1.

The disease status is split into disease specific sections:

- Acute leukemias
- Chronic leukemias
- Plasma cell neoplasms
- MPN, MDS, MDS/MPN overlap syndromes
- Lymphomas
- Solid tumours
- Bone marrow failure syndromes (BMF) including aplastic anaemia (AA)
- Autoimmune disorders
- Haemoglobinopathies
- Other diagnoses
- Inborn errors

The instructions for completing appendix 1 can be found in appendix 1 of this document.

Pregnancy after HCT

Has the patient become pregnant or impregnated another person since the last follow-up?

If the patient has not become pregnant or has not impregnated another person since the last follow-up, select **No** and proceed to the next section.

Extended dataset

Was there an attempted pregnancy since the last follow-up?

Indicate if there was an attempted pregnancy since the last follow-up select **Yes, No or Unknown.**

If the patient has become pregnant or has impregnated another person since the last follow-up select **Yes** and provide details. Select Unknown if it is not known.

Did the pregnancy result in a live birth?

If the patient has become pregnant or has impregnated another person since the last follow-up, answer **Yes**. If the pregnancy resulted in a live birth, indicate the **Year of birth** and **Month of birth** of the child, or mark the date **Unknown** if the date is not available.

Answer **No** if pregnancy did not result in a live birth and indicate the **Date of spontaneous or induced termination** (YYYY/MM/DD) or mark the date as **Unknown**. In case of multiple spontaneous or induced terminations, report the date of the first such event.

Select **Still pregnant at time of follow-up** if the patient/the person they impregnated was still pregnant at the time of the follow-up.

Extended dataset

Conception method

Indicate which conception method was used for the pregnancy that resulted in live birth. Select **Natural, Assisted** or **Unknown**

If there is no detailed information about the pregnancy and whether or not it resulted in a live birth, select **Unknown**.

If multiple pregnancies occurred in the follow-up period, the live birth should be prioritised in reporting, in the absence of the live birth, ongoing pregnancy should be prioritised: if one pregnancy resulted in live birth and another not, report the live birth only; if there occurred any terminated pregnancy and the other pregnancy is ongoing as of this follow-up date, select *Still pregnant at time of follow-up* answer option.

Appendix 1 - Disease specific best response and disease status

The disease status and best response are split into disease specific sections which can be found in appendix 1 of the follow-up form. This section is separated into disease status for:

- Acute leukaemias;
- Chronic leukemias;
- Plasma cell neoplasms;
- MPN, MDS, MDS/MPN overlap syndromes;
- Lymphomas;
- Solid tumours;
- Bone marrow failure syndromes (BMF) including aplastic anaemia (AA);
- Autoimmune disorders;
- Haemoglobinopathies;
- Other diagnoses;
- Inborn errors .

Please make sure to check the **Disease specific best response and disease status v1** document latest version available under *Manuals and Reference Documents* section on [EBMT website](#).

Acute leukaemias

This section is applicable to acute myeloid leukaemias (AML), precursor lymphoid neoplasms (PLN) and other acute leukaemias.

Acute leukaemias disease status or best response

Select the disease status or best response from the following list, refer from the criteria described in the **Disease specific best response and disease status v1** document.

- Complete remission (CR)
- Not in complete remission

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively. If reporting the disease status, also indicate the minimal residual disease (MRD) status.

Chronic leukaemias

The chronic leukaemias section is split into chronic myeloid leukaemia (CML) and chronic lymphocytic leukaemia (CLL), prolymphocytic leukaemia (PLL) and other chronic leukaemias.

Chronic myeloid leukaemia disease status or best response

Select the disease status or best response from the list:

- Chronic phase (CP) and type of remission (haematological, cytogenetic, molecular)
- Accelerated phase
- Blast crisis

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Number

If the disease status or best response was chronic phase (CP), accelerated phase (AP) or blast crisis (BC), select the number of this status.

Number the different disease statuses chronologically. A patient can only be in the next chronic phase after he has experienced a blast crisis or accelerated phase.

If the disease status or best response was chronic phase (CP) also indicate:

Haematological remission

If the patient was in Chronic phase (CP), report if haematological remission was achieved (answer Yes), or not achieved (answer No) according to the criteria provided in the **Disease specific best response and disease status** document . Answer **Not evaluated** if it was not evaluated or Unknown if it cannot be verified if it was evaluated or not.

Cytogenetic remission

If the patient was in Chronic phase (CP), report if cytogenetic remission was achieved (answer Yes), or not achieved (answer No) according to the criteria provided in the **Disease specific best response and disease status** document. Answer Not evaluated if it was not evaluated or Unknown if it cannot be verified if it was evaluated or not.

Note: A patient in cytogenetic remission must be in haematological remission but could still present a molecular relapse. This is because the cytogenetic technique has a higher resolution than haematological measurements but lower resolution than molecular methods.

*Extended dataset**Cytogenetic detection*

Please answer these questions when the patient is in accelerated phase, blast crisis, or chronic phase without cytogenetic remission.

t(9;22) positive metaphases

Please report here the percentage (%) of the metaphases with the t(9;22) translocation. If the test was not done, select **not evaluated**. If the value is not known, select **unknown**.

t(9;22) positive cells detected by FISH

FISH (Fluorescent In Situ Hybridisation) is another cytogenetic technique that is frequently used. Where “conventional” cytogenetic investigation is performed on dividing cells (metaphases), FISH can also analyse non-dividing bone marrow cells (cells in interphase) and investigates about 400 cells, thus many more than the cells that are analysed with “conventional” cytogenetic analysis.

Molecular remission

If the patient was in Chronic phase (CP), report if molecular remission was achieved (answer Yes), or not achieved (answer No) according to the criteria provided in the **Disease specific best response and disease status** document. Answer Not evaluated if it was not evaluated or Unknown if it cannot be verified if it was evaluated or not.

Note: A patient in molecular remission must also be in cytogenetic and haematological remission. This is because molecular techniques have a higher resolution than both haematological and cytogenetic measurements.

*Extended dataset**BCR::ABL1 variant allele frequency (VAF)*

Please answer this question when the patient is in an accelerated phase, blast crisis, or chronic phase without molecular remission.

Please report the variant allele frequency of BCR::ABL1.

Chronic lymphocytic leukaemia (CLL), prolymphocytic leukaemia (PLL) and other chronic leukaemias disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status v1** document:

- Complete Remission (CR)
- Partial Remission (PR)
- Progression
- Stable Disease (SD)
- Relapse
 - Relapse is defined as evidence of disease progression in a patient who has previously achieved the criteria of a CR or PR for ≥ 6 months : see the **Disease specific best response and disease status** document.

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Progression sensitivity

If the disease status or best response was progression, indicate if the progression was **resistant** to the last chemotherapy regimen the patient received, or if it was **sensitive**. If this is not known, select **unknown**.

Plasma cell neoplasms

Disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status** document:

- Complete remission (CR)
- Stringent complete remission (sCR)
- Very good partial remission (VGPR)
- Partial remission (PR)
- Stable disease (no change, no response/loss of response)
- Progression
- Relapse

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Please check the **Disease specific best response and disease status v1** document with the criteria for each disease status or response category for plasma cell neoplasms and a criteria for Immunoglobulin-related (AL) Amyloidosis.

Number

Please report the status/response number (if it is **1st, 2nd, 3rd or higher** or if it is **Unknown**) for the following disease statuses or responses:

- Complete remission (CR);
- Stringent complete remission (sCR);
- Very good partial remission (VGPR);
- Partial remission (PR);
- Relapse.

Each different status/response has their own sequential count.

For example, a patient received a non-graft treatment and is in PR1 as response to this treatment. After that there is a progression, another treatment, and response to this treatment is CR1, relapse, another treatment, response CR2.

The count doesn't reflect the different disease statuses/responses (eg. in the example above it should not be PR1, CR2, CR3), but within that status/response the sequential count (so PR1, CR1, CR2).

Extended dataset

Immunoglobulin-related (AL) amyloidosis

Organ response

Please check the **Disease specific best response and disease status v1** document for definitions of organ response and progression.

Was the patient on dialysis during this follow-up period?

Report whether the patient was on dialysis during this follow-up period. Select **Unknown** if this information is unavailable.

Started in this follow-up period

Select this option if dialysis started during this follow-up period.

Start date

If the answer to the previous question was **Yes**, report the start date of dialysis. If the start date is not known, select **Unknown**.

Ongoing since previous follow-up

Select this option if dialysis started during a previous follow-up period and was still ongoing during this follow-up.

Did dialysis stop?

Report whether dialysis was stopped during this follow-up. Select **Unknown** if this information is unavailable.

End date

Report the dialysis end date. If the dialysis stopped but the end date is not known, select **Unknown**.

Minimal residual disease

Complete this section only if the patient disease status is reported for patients with an AL, CLL leukaemia or a plasma cell neoplasm diagnosis.

If the patient is in CR or sCR, but has evidence of disease by more sensitive assessments including molecular, flow cytometry or cytogenetic methods, mark it as **Positive**. If the MRD assay cannot detect malignant cells mark it as **Negative**. Mark it as **Not evaluated** if MRD status evaluation was not carried out during the follow-up period.

Date MRD status evaluated

Report the date the MRD status was evaluated. If the date is not known, select **unknown**.

Extended dataset

Sensitivity of MRD assay

Select the appropriate sensitivity of the MRD assay from the list.

Method used

Select the most sensitive method that was used to assess the MRD status.

Myeloproliferative neoplasms (MPN), Myelodysplastic neoplasms (MDS), MDS/MPN overlap syndromes

Disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status v1** document:

- Complete remission (CR);
- Improvement but no CR;
- Primary refractory phase (no change);
- Relapse;
- Progression/worsening.

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Number

If the disease status or best response was complete remission (CR) or relapse, please report the number (if it is **1st, 2nd, 3rd or higher** or if it is **Unknown**).

Each different status/response has their own sequential count.

For example, a patient received a non-graft treatment and is CR1 in response to this treatment, after that there is a (1st) relapse, another treatment, and response CR2.

The count doesn't reflect the different disease statuses/responses (eg. in the example above it should not be CR1, 2nd Relapse, CR3), but within that status/response the sequential count (so CR1, 1st Relapse, CR2).

Autoimmune disorders

Disease status or best response

Select the disease status or best response from the list:

- **No evidence of disease**- the patient has achieved complete absence of disease, there are no signs or symptoms of the original disease described.
- **Improved**
- **Unchanged** - Patients who have not demonstrated complete absence of disease, improvement in symptoms, or deterioration of symptoms will be classified as **Unchanged**.
- **Worse**

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Haemoglobinopathies

The haemoglobinopathies section is split into thalassemia and sickle cell disease.

Thalassemia best response

Select the best response that was achieved since the main treatment:

- Transfusion independent;
- Transfusions required.

For clarification, transfusion independence is typically defined as going 8-12 weeks without needing transfusions, without a specific haemoglobin threshold.

If the best response is not known or was not evaluated, select **Unknown** or **Not evaluated**, respectively.

Date of last transfusion

If a patient reached transfusion independence, report the date of the last transfusion after main treatment the patient received. If the date is not known, select **Unknown**.

Date of first transfusion

If a patient still requires transfusions, report the date of the first transfusion the patient received after main treatment due to haemoglobin deficiency (recurrence of disease). If the date is not known, select **Unknown**.

Thalassemia disease status

Patient requires transfusions during follow-up period

Indicate if the patient requires transfusions during follow-up period after haematopoietic recovery by selecting **No** or **Yes**. Mark as Unknown if it is not known.

Patient requires transfusions, Yes

If a patient is transfusion dependent indicate whether at this follow-up period the **Return to transfusion dependence after HCT or transfusion free period** occurred or patient **Ongoing transfusion dependence since previous assessment**.

Date of first transfusion

If a patient has returned to transfusion dependence after main treatment or transfusion free period during this follow-up period, report the date of the first transfusion after main treatment or transfusion free period. If the date is not known, select **Unknown**. If a patient has ongoing transfusion dependence since the previous assessment and the date was reported at the previous follow-up form, do not report the date here, the question will be disabled.

Number of units

Report the number of transfusion units patient received during this follow-up period. If the exact number is not known, select **Unknown**.

Did transfusions stop?

Indicate if the patient stopped receiving the transfusions. If the patient stopped transfusions and did not require more during this follow-up period, select **Yes**.

Date of last transfusion

If the patient stopped transfusions during the reporting period, provide the date when the last transfusion was administered. If it is not known, select **Unknown**.

Sickle cell disease best response

Select the best response that was achieved since the main treatment:

- No return of sickling episodes. Patients are considered to have no return of sickling episodes when they have shown an absence of sickle cell crises.
- Return of sickling episodes. When sickle cell crises reoccur.

If the best response is not known or was not evaluated, select **Unknown** or **Not evaluated**, respectively.

Date of first episode

If a patient has returned to sickling episodes, report the date of the first episode after main treatment. If the date is not known, select **unknown**.

Sickling episodes occur during follow-up period

This should only be completed when reporting the disease status. Indicate if there were no more recurrent sickle cell episodes after HCT by selecting **No**. If recurrent sickling episodes were present after HCT, select **Yes**. If it is not known if the sickling episodes returned, select **Unknown**.

Sickling episodes occur during follow-up period, Yes

Indicate whether the sickling episodes first return after HCT or there was an ongoing presence of sickling episodes since last follow-up assessment.

Date of first episode

If the sickling episodes reoccurred for the first time since the main treatment, report the date of the first sickling episode. If the date is not known, select **Unknown**.

Number of SCD episodes (during follow-up)

If the sickling episodes first returned after main treatment, report the number of sickling episodes which occurred during this follow-up period. If the number is not known, select **Unknown**.

Lymphomas

Disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status** document:

- Chemorefractory relapse or progression, including primary refractory disease
- Complete remission (CR)
- Partial remission
- Stable disease (no change, no response/loss of response)
- Untreated relapse (from a previous CR) or progression (from a previous PR)

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Technique used for disease assessment

Specify the technique used to assess the disease status or best response.

If the technique used is not listed, select **Other** and specify the technique used in the text field.

Solid tumours

Disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status v1** document:

- Complete remission (CR)
- First partial remission
- Partial remission (PR)
- Progressive disease
- Relapse
- Stable disease (no change, no response/loss of response)

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Bone marrow failures (incl. AA)

Disease status or best response

Select the disease status or best response from the list according to the criteria provided in the **Disease specific best response and disease status** document:

- Complete remission (CR)
- Partial remission (PR)
- Haematological improvement (HI); NIH partial response
- Stable disease (no change, no response/loss of response)
- Relapse/Progression

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Did transfusions stop during the follow-up period?

If the disease status is reported for bone marrow failures, indicate if transfusions stopped since the last follow-up. If the patient was never transfusion dependent, select **patient was never transfusion dependent**.

If the transfusions are ongoing, select **no**. If the transfusions did stop, select **Yes** and complete the next questions. If the patient is transfusion independent and was transfusion dependent at a previous follow-up, select **ongoing transfusion independence since last follow-up**. If the transfusion status is not known, select **unknown**.

Did the patient return to transfusion dependency afterwards?

If the patient became transfusion independent since the last follow-up, but is back to needing transfusions within this follow-up period, select **yes**. If they continue to be independent of transfusions after stopping in this follow-up period, select **no**. If it is not known if the patient went back to needing transfusions, select **unknown**.

First transfusion date

If the patient stopped transfusions during the follow-up period but went back to being transfusion dependent, report the first transfusion date after the transfusion free period. If the date is not known, select **unknown**.

Other diagnosis

Disease status or best response

Select the disease status or best response from the list:

- **No evidence of disease** - the patient has achieved complete absence of disease, there are no signs or symptoms of the original disease described.
- **Improved**
- **No response**
- **Worse**

If the disease status or best response is not known or was not evaluated, select **unknown** or **not evaluated**, respectively.

Disease status (Inborn errors only)

Patient height at this follow-up

Report the height of the patient at the last follow-up in centimetres.

Patient weight at this follow-up

Report the weight of the patient at the last follow-up in kilograms.

Patient is attending

Please select whether one of the situation below applies to the patient:

- **Regular school/work:** If the patient attending regular school or work;
- **Alternative school/adapted work:** If the patient attending school or working that is adapted to their inabilities;
- **Patient is not able to attend work/school:** if the patient does not attend school or work

If the patient is not known whether he/she is attending school or working, please select **Unknown**.

Immune profiling done during this follow-up period

Only for Inborn errors of Immunity: Indicate if immune profiling was done during this follow-up period.

Test date

Report the date of the most recent immune reconstitution was tested

Cell type and test results

Please report the value of the cell types below from the most recent test performed. For Naive CD4 T-cells and CD8 T-cells please select the unit of measurements.

Cell type	Unit of measurement
CD3 T-cells (CD3)	Cells/ μ l
CD4 T-cells (CD4)	Cells/ μ l
CD8 T-cells (CD8)	Cells/ μ l
B-cells (CD19)	Cells/ μ l
NK-cells (CD16/CD56)	Cells/ μ l
Naive CD4 T-cells (CD4/CD45RA)	Please select the unit of measurement
Naive CD8 T-cells (CD8/CD45RA)	Please select the unit of measurement
IgG	Gram/l
IgA	Gram/l
IgM	Gram/l

Extended dataset

Disease status (Inborn errors only)

Select the immunomodulatory treatments the patient received in the 3 months before the follow-up

Only report treatments administered in the 3 months before this follow-up for the underlying disease. Do not report treatments for GvHD or other HCT/CT related complications.

Comorbidities during this follow up period

Inborn errors of Immunity only: indicate if the comorbidities de novo, resolved, improved, stabilised or worsened since the last follow-up.

Was the patient admitted to ICU during this follow-up period

Indicate if the patient was admitted to ICU during this follow-up period. If the information is not available mark as **Unknown**.

Cell Infusion Sheet

The following completion guidelines refer to the completion of appendix 6 of the annual/unscheduled follow-up form, the cell infusion sheet.

Please report each cell infusion episode performed during the follow-up period after HCT in a separate cell infusion sheet, completing as many sheets as episodes of cell infusion that took place. It is important that cell infusions for the prevention or treatment of complications are reported here and not on the Cell Therapy forms. Cell infusion treatment is often given as sequential cell infusions through a series of days or even weeks. In order to make the data comparable, one episode of cell infusion treatment (one "CI") is defined as any number of cell infusions that take place for the same indication within 10 weeks from first to last infusion. If the indication for the treatment changes within the 10 weeks, that would be considered as 2 separate episodes of cell infusion (2 "CI"), with the 2nd episode starting on the 1st day infusions were given after the change in indication.

Do not use this cell infusion sheet for any boost. All boosts shall be registered inside the HCT follow-up form.

Chronological number of CI episode for this patient

Report the chronological number of this cell infusion episode for this patient.

Date of the first infusion

Report the date of the first infusion within this episode.

Number of infusions within 10 weeks

Report the number of infusions within 10 weeks. Count only infusions that are part of the same regimen and given for the same indication.

Source of cells

Indicate if the source of cells are **allogeneic** and/or **autologous**, check all that apply.

Type of cells

Select the type of cells, check all that apply:

- **Lymphocytes (DLI);**
- **Mesenchymal;**
- **Fibroblasts;**
- **Dendritic cells;**
- **NK cells;**
- **Regulatory T-cells;**

- **Gamma/delta cells;**
- **Virus-specific T-cells (VST).**

If the type of cells is not listed, select **Other** and specify the type of cells in the text field in English.

Specificity of VST product

If the type of cells is '**Virus-specific T-cells (VST)**', indicate whether VST were '**Single-virus specific**' or '**Multi-virus specific**'.

If single-virus specific, to which virus?

If '**Single-virus specific**', check to which virus the VST are specific. Please register this virus in the Infectious complications - Viral infection part and fill in the VST in the 'Pre-emptive viral therapy' or 'Treatment of end-organ viral disease' section.

If multi-virus specific, to which viruses?

If '**Multi-virus specific**', check all viruses to which the VST are specific. Please register these viruses in the Infectious complications - Viral infection part and fill in the VST in the 'Pre-emptive viral therapy' or 'Treatment of end-organ viral disease' section.

Extended dataset

Product origin

If the type of cells is '**Virus-specific T-cells (VST)**', indicate whether the VST product was '**In-house manufactured**' or is a '**Commercial product**'.

If the VST product is a '**Commercial product**' check the box if this commercial product is '**Tabelecleucel**'.

Donor source

If the source of cells is '**Allogeneic**', indicate whether this source was the '**Stem cell donor**' or a '**Third-party donor**'.

If the source was a '**Third party donor**', indicate whether this donor was '**Related (family)**' or '**Unrelated**'.

Manufacturing method

Indicate the method by which the VST were manufactured.

Number of doses per infection episode

Indicate whether the patient received 1 VST dose or more than 1 VST dose per infection episode.

Total number of doses

If the patient received more than 1 VST dose per infection episode, indicate how many doses the patient received in total.

Dosing interval (in days)

If the patient received more than 1 VST dose per infection episode, indicate how many days were in between 2 consecutive doses.

Disease status at time of this cell infusion

Indicate the disease status at time of this cell infusion corresponding to indication diagnosis by selecting from the list provided in Appendix 1.

The disease status is split into disease specific sections:

- Acute leukemias;
- Chronic leukemias;
- Plasma cell neoplasms;
- MPN, MDS, MDS/MPN overlap syndromes;
- Lymphomas;
- Solid tumours;
- Bone marrow failure syndromes (BMF) including aplastic anaemia (AA);
- Autoimmune disorders;
- Haemoglobinopathies;
- Other diagnoses.

The instructions for completing appendix 1 can be found in appendix 1 of this document.

If the disease status has not been evaluated, select **Not evaluated**. Select **Unknown** if the disease status at the time of this cell infusion is not known.

Indication

Select all the indications for this cell infusion episode that apply:

- **Planned/protocol;**
- **Prophylactic;**
- **Treatment of acute GvHD;**
- **Treatment of chronic GvHD;**
- **Treatment PTLD, EBV lymphoma;**
- **Treatment for primary disease;**

- **Mixed chimaerism;**
- **Loss/decreased chimaerism;**
- **Treatment of viral infection other than EBV;**
- **Poor graft function;**
- **Infection prophylaxis.**

If the indication is not listed, select **Other** and specify it in the text field in English.

Response to DLI:

Only for Acute leukemia donor lymphocyte infusions: specify the response to DLI or mark it as Unknown if it is not known.

MRD status

If the response to DLI is reported as Complete remission (CR), report also the MRD status: if it is MRD negative or positive. Select **Not evaluated** if it was not evaluated or **Unknown** if it is not known.

Bibliography

1. Jagasia MH, Greinix HT, Arora M, Williams KM, Wolff D, Cowen EW, et al. National Institutes of Health Consensus Development Project on Criteria for Clinical Trials in Chronic Graft-versus-Host Disease: I. The 2014 Diagnosis and Staging Working Group report. *Biol Blood Marrow Transplant* [Internet]. 2015 Mar;21(3):389–401.e1. Available from: <http://dx.doi.org/10.1016/j.bbmt.2014.12.001>
2. Schoemans HM, Lee SJ, Ferrara JL, Wolff D, Levine JE, Schultz KR, et al. EBMT-NIH-CIBMTR Task Force position statement on standardized terminology & guidance for graft-versus-host disease assessment. *Bone Marrow Transplant* [Internet]. 2018 Nov;53(11):1401–15. Available from: <http://dx.doi.org/10.1038/s41409-018-0204-7>
3. Holler E, Greinix H, Zeiser R. Acute Graft-Versus-Host Disease. In: Carreras E, Dufour C, Mohty M, Kröger N, editors. *The EBMT Handbook: Hematopoietic Stem Cell Transplantation and Cellular Therapies* [Internet]. Cham (CH): Springer; Available from: http://dx.doi.org/10.1007/978-3-030-02278-5_43
4. Wolff D, Lawitschka A. Chronic Graft-Versus-Host Disease. In: Carreras E, Dufour C, Mohty M, Kröger N, editors. *The EBMT Handbook: Hematopoietic Stem Cell Transplantation and Cellular Therapies* [Internet]. Cham (CH): Springer; Available from: http://dx.doi.org/10.1007/978-3-030-02278-5_44
5. Cuvelier GDE, Schoettler M, Buxbaum NP, Pinal-Fernandez I, Schmalzing M, Distler JHW, et al. Toward a Better Understanding of the Atypical Features of Chronic Graft-Versus-Host Disease: A Report from the 2020 National Institutes of Health Consensus Project Task Force. *Transplant Cell Ther* [Internet]. 2022 Aug;28(8):426–45. Available from: <http://dx.doi.org/10.1016/j.jtct.2022.05.038>
6. National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE) [Internet]. 2021 [cited 2023 Mar 21]. Available from: https://ctep.cancer.gov/protocoldevelopment/electronic_applications/ctc.htm
7. Schoettler ML, Carreras E, Cho B, Dandoy CE, Ho VT, Jodele S, et al. Harmonizing Definitions for Diagnostic Criteria and Prognostic Assessment of Transplantation-Associated Thrombotic Microangiopathy: A Report on Behalf of the European Society for Blood and Marrow Transplantation, American Society for Transplantation and Cellular Therapy, Asia-Pacific Blood and Marrow Transplantation Group, and Center for International Blood and Marrow Transplant Research. *Transplant Cell Ther* [Internet]. 2023 Mar;29(3):151–63. Available from: <http://dx.doi.org/10.1016/j.jtct.2022.11.015>

8. Ruutu T, Carreras E. Hepatic Complications. In: Carreras E, Dufour C, Mohty M, Kröger N, editors. The EBMT Handbook: Hematopoietic Stem Cell Transplantation and Cellular Therapies [Internet]. Cham (CH): Springer; Available from: http://dx.doi.org/10.1007/978-3-030-02278-5_49
9. Mermel LA, Allon M, Bouza E, Craven DE, Flynn P, O'Grady NP, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 Update by the Infectious Diseases Society of America. Clin Infect Dis [Internet]. 2009 Jul 1;49(1):1–45. Available from: <http://dx.doi.org/10.1086/599376>
10. Comenzo RL, Reece D, Palladini G, et al. Consensus guidelines for the conduct and reporting of clinical trials in systemic light-chain amyloidosis. Leukemia 2012; 26:2317.